

SALZIX WART REMOVER- salicylic acid liquid
Toprail Chp Llc

SALZIX LIQUID WART REMOVER

Drug Facts

Active ingredient

Salicylic acid 17% w/w

Purpose

Wart remover

Use

- for the removal of common warts and plantar warts on the bottom of the foot. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness, and the interruption of the footprint pattern.

Warnings

For external use only.

Flammable • keep away from fire or flame

Do not use • if you are diabetic or if you have poor blood circulation

- on genital warts or warts on the face or mucous membrane
- on moles, birthmarks or warts with hair growing from them
- on irritated skin or on any area that is infected or reddened

When using this product • If product gets into the eye, flush with water for 15 minutes • avoid inhaling vapors • **cap bottle tightly** and store at room temperature away from heat

Stop use and ask a doctor if discomfort persists.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- wash affected area, may soak wart in warm water for 5 minutes • dry area thoroughly
- carefully apply one drop at a time with applicator to sufficiently cover each wart • let dry • repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks

Other information

• store between 20°-25°C (68°-77°F) • **close tightly to prevent solidification** • **wipe top of bottle clean after each use** • discard when signs of wart no longer appear • avoid surrounding skin when applying product • do not allow to come into contact with fabrics, floors, countertops or other surfaces - will stain

Inactive ingredients

camphor, castor oil, ethanol, ethyl ether, nitrocellulose

Questions?

Call 1-866-964-0939

- **Removes Common & Plantar Warts**
- **Convenient Brush Applicator**
- **Safe & Effective**
- **Maximum Strength**
- **Fast Acting Liquid**

IMPORTANT: CLOSE CAP TIGHTLY AFTER USE

DO NOT USE IF NECKBAND IS BROKEN

DISTRIBUTED BY:

TOPRAIL PHARMACEUTICALS

NORTHBROOK, IL

www.TOPRAILPHARMACEUTICALS.com

MADE IN USA FROM GLOBALLY SOURCED MATERIALS

Keep packaging for complete drug facts information.

Packaging

Outer Package Label



- Removes Common & Plantar Warts
- Convenient Brush Applicator
- Safe & Effective
- Maximum Strength
- Fast Acting Liquid




Not Actual Size

NDC# 72847-333-29 NET 0.31 FL. OZ. (9 mL)

- Removes common & plantar warts
- Convenient brush applicator
- Safe & effective

Use for warts on feet & hands



53-012TR

TOPRAIL
PHARMACEUTICALS

MAXIMUM STRENGTH

LIQUID WART REMOVER

Salicylic Acid 17% / Wart Remover

NDC# 72847-333-29 NET 0.31 FL. OZ. (9 mL)

**NO VARNISH
AREA
(WHITE ONLY)**

DISTRIBUTED BY:
TOPRAIL PHARMACEUTICALS
NORTHBROOK, IL
www.TOPRAILPHARMACEUTICALS.com



NDC# 72847-333-29
MADE IN USA FROM
GLOBALLY SOURCED MATERIALS

Drug Facts

Active ingredient Salicylic acid 17% w/w. Wart remover

Purpose

Use ■ for the removal of common warts and plantar warts on the bottom of the foot. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness, and the interruption of the footprint pattern.

Warnings

For external use only.

Flammable ■ keep away from fire or flame

Do not use ■ if you are diabetic or if you have poor blood circulation

■ on genital warts or warts on the face or mucous membrane

■ on moles, birthmarks or warts with hair growing from them

■ on irritated skin or on any area that is infected or reddened

When using this product ■ if product gets into the eye, flush with water for 15 minutes ■ avoid inhaling vapors ■ cap bottle tightly

Stop use and ask a doctor if discomfort persists.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ wash affected area, may soak wart in warm water for 5 minutes ■ dry area thoroughly ■ carefully apply one drop at a time with applicator to sufficiently cover each wart ■ let dry ■ repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks

Other information ■ store between 20°–25°C (68°–77°F) ■ close tightly to prevent solidification ■ wipe top of bottle clean after each use ■ discard when signs of wart no longer appear ■ avoid surrounding skin when applying product ■ do not allow to come into contact with fabrics, floors, countertops or other surfaces — will stain

Inactive ingredients camphor, castor oil, ethanol, ethyl ether, nitrocellulose

Questions? Call 1-866-964-0939



Inner Package Label

DO NOT USE IF NECKBAND IS BROKEN

NO VARNISH AREA
50-012TR0.5315"

SALZIX
Salicylic Acid 17% / Wart Remover

LIQUID WART REMOVER

- Removes Common & Plantar Warts
- Convenient Brush Applicator
- Safe & Effective

CLOSE CAP TIGHTLY AFTER USE
MAXIMUM STRENGTH
NDC# 72847-333-29
NET 0.31 FL. OZ. (9 mL)

Active ingredient: Salicylic acid (17% w/w) **Purpose:** Wart remover **Use:** For the removal of common warts and plantar warts on the bottom of the foot. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness and the interruption of the footprint pattern. **Warnings:** For external use only. **Flammable:** Keep away from fire or flame.

Do not use if you are diabetic or if you have poor blood circulation, on genital warts or warts on the face or mucous membranes, on moles, birthmarks or warts with hair growing from them, or on irritated skin or on any area that is infected or reddened. **When using this product** if product gets into the eye, flush with water for 15 minutes. Avoid inhaling vapors. **Cap bottle tightly** and store at room temperature away from heat. **Stop use and ask a doctor** if discomfort persists. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away. **Directions:** Wash affected area thoroughly. Carefully apply one drop at a time with applicator to sufficiently cover each wart. Let dry. Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks. **Other information:** Store between 20° – 25°C (68° – 77°F). **Close tightly to prevent solidification. Wipe top of bottle clean after each use.** Discard when signs of wart no longer appear. Avoid surrounding skin when applying product. Do not allow to come into contact with fabrics, floors, countertops or other surfaces -- will stain. **Inactive ingredients:** camphor, castor oil, ethanol, ethyl ether, nitrocellulose **Questions?** Call 1-866-964-0939 **Keep packaging for complete drug facts information.** Made in USA from globally sourced materials. Distributed by: TOPPALL PHARMACEUTICALS, NORTHBROOK, IL

Drug Facts Label

Drug Facts

Active ingredient	Purpose
Salicylic acid 17% w/w.....	Wart remover

Use ■ for the removal of common warts and plantar warts on the bottom of the foot. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness, and the interruption of the footprint pattern.

Warnings

For external use only.

Flammable ■ keep away from fire or flame

Do not use ■ if you are diabetic or if you have poor blood circulation
 ■ on genital warts or warts on the face or mucous membrane
 ■ on moles, birthmarks or warts with hair growing from them
 ■ on irritated skin or on any area that is infected or reddened

When using this product ■ If product gets into the eye, flush with water for 15 minutes ■ avoid inhaling vapors ■ **cap bottle tightly** and store at room temperature away from heat

Stop use and ask a doctor if discomfort persists.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ wash affected area, may soak wart in warm water for 5 minutes ■ dry area thoroughly ■ carefully apply one drop at a time with applicator to sufficiently cover each wart ■ let dry ■ repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks

Other information ■ store between 20° – 25°C (68° – 77°F)
 ■ **close tightly to prevent solidification** ■ **wipe top of bottle clean after each use** ■ discard when signs of wart no longer appear

after each use ■ discard when signs of wart no longer appear
■ avoid surrounding skin when applying product ■ do not allow to come
into contact with fabrics, floors, countertops or other surfaces -- will stain

Inactive ingredients camphor, castor oil, ethanol, ethyl ether,
nitrocellulose

Questions? Call 1-866-964-0939

SALZIX WART REMOVER

salicylic acid liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72847-333
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	0.17 g in 9 mL

Inactive Ingredients

Ingredient Name	Strength
CAMPHOR (NATURAL) (UNII: N20HL7Q941)	
CASTOR OIL (UNII: D5340Y2I9G)	
ALCOHOL (UNII: 3K9958V90M)	
ETHER (UNII: 0F5N573A2Y)	
PYROXYLIN (UNII: KYR8BR2X6O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72847-333-29	1 in 1 BOX	03/20/2025	
1		9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	03/20/2025	

Labeler - Toprail Chp Llc (080783804)