

DAYTIME COLD AND FLU- daytime cold and flu capsule, liquid filled

Allegiant Health

474 - Daytime Cold and Flu Relief

Active ingredient(s)

Acetaminophen 325mg

Dextromethorphan HBr 10mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Use(s)

temporarily relieves these symptoms due to cold flu:

- cough
- sore throat
- headache
- minor aches and pains
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 10 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use if you have

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if

if you are taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breastfeeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222)

right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose

Adults and children 12 years and over: Take 2 softgels with water every 4

hours. Do not exceed 8 softgels in 24 hours or as directed by a doctor

Children under 12 years: Do not use

Other information

- store at room temperature 20-25°C (68-77°F)
- avoid excessive heat
- do not use if blister unit is torn or open

Inactive ingredients

FD&C Red #40, FD&C Yellow #6, Gelatin, Glycerin, Polyethylene Glycol 400, Povidone K12, Propylene Glycol, Purified water, Sorbitol Sorbitan Solution, White Edible Ink

Questions/Comments

Call **1-888-952-0050**

(Monday-Friday 9 AM – 5 PM EST)

Principal Display Panel

Drug Facts**Active ingredients (in each softgel)**

Dextromethorphan HBr 10mg.....Cough suppressant

Uses temporarily relieves these symptoms due to a cold or flu:
 in cough in sore throat in headache in minor aches and pains
 in temporarily reduces fever

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take in more than 10 softgels in 24 hours, which is the maximum daily amount for this product. in with other drugs containing acetaminophen in 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: in skin reddening in blisters in rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use in with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 in if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. in if you have ever had an allergic reaction to this product or any of its ingredients in in children under 12 years of age.

Ask a doctor before use if you have: in liver disease in heart disease in thyroid disease in diabetes in high blood pressure in difficulty in urination due to enlargement of the prostate gland in

Drug Facts (continued)

in cough with excessive phlegm (mucus) in persistent or chronic cough as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if in nervousness, dizziness or sleeplessness occurs in pain, cough gets worse or lasts more than 7 days in fever gets worse or lasts more than 3 days in redness or swelling is present in new symptoms occur in cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions in do not take more than the recommended dose.
Adults and children 12 years and over Take 2 softgels with water every 4 hours. Do not exceed 8 softgels in 24 hours or as directed by a doctor.

Children under 12 years

Do not use

Other information in store at room temperature 20-25°C (68-77°F) in avoid excessive heat in do not use if blister unit is torn or open.

Inactive ingredients: FD&C Red #40, FD&C Yellow #6, Gelatin, Glycerin, Polyethylene Glycol 400, Povidone K12, Propylene Glycol, Purified water, Sorbitol Sorbitan Solution, White Edible Ink.

Questions or comments? Call 1-888-652-0090 Monday through Friday 9AM - 5PM EST.

HealthAzz® DAYTIME

COLD & FLU RELIEF

Relieves:
 Cough Sore Throat Headache
 Minor Aches & Pains
 Temporarily Reduces Fever

20 Softgels



Acetaminophen 325mg / Pain Reliever / Fever Reducer
 Dextromethorphan HBr 10mg / Cough Suppressant

Safe For Adults With
 High Blood Pressure

DAYTIME

HealthAzz®

NO OTHER DRUGS
 CONTAINING
 ACETAMINOPHEN
 OR
 DEXTROMETHORPHAN
 SHOULD BE TAKEN
 WITH THIS PRODUCT

Safe For Adults With
 High Blood Pressure

DAYTIME

DAYTIME COLD AND FLU

daytime cold and flu capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-474
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K12 (UNII: 333AG72FWJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	P41
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-474-29	1 in 1 CARTON	01/09/2026	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69168-474-51	2 in 1 CARTON	01/09/2026	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/09/2026	

Labeler - Allegiant Health (079501930)

Revised: 1/2026

Allegiant Health