
CEFOXITIN FOR INJECTION, USP

PHARMACY BULK PACKAGE NOT FOR DIRECT INFUSION

Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefoxitin for Injection, USP and other antibacterial drugs, Cefoxitin for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Cefoxitin for Injection, USP is a semi-synthetic, broad-spectrum cepha antibiotic sealed under nitrogen for intravenous administration. It is derived from cephamycin C, which is produced by *Streptomyces lactamdurans*. Its chemical name is sodium (6R,7S)-3-(hydroxymethyl)-7-methoxy-8-oxo-7-[2-(2-thienyl)acetamido]-5-thia-1-azabicyclo [4.2.0] oct-2-ene-2-carboxylate carbamate (ester).

The molecular formula is $C_{16}H_{16}N_3N_aO_7S_2$, and the structural formula is:

Cefoxitin for injection contains approximately 53.8 mg (2.3 milliequivalents) of sodium per gram of cefoxitin activity. Solutions of Cefoxitin for Injection, USP range from colorless to light amber in color. The pH of freshly constituted solutions usually ranges from 4.2 to 7.0.

Each Pharmacy Bulk Package Bottle of Cefoxitin for Injection, USP contains sterile cefoxitin sodium, USP equivalent to 10 grams of cefoxitin.

This Pharmacy Bulk Package Bottle is a container of sterile preparation for intravenous use that contains many single doses. The contents are intended for use in a pharmacy admixture service and are restricted to the preparation of admixtures for intravenous infusion. AFTER RECONSTITUTION FURTHER DILUTION IS REQUIRED BEFORE USE. NOT FOR DIRECT INFUSION.

CLINICAL PHARMACOLOGY

Clinical Pharmacology

Following an intravenous dose of 1 gram, serum concentrations were 110 mcg/mL at 5 minutes, declining to less than 1 mcg/mL at 4 hours. The half-life after an intravenous dose is 41 to 59 minutes. Approximately 85 percent of cefoxitin is excreted unchanged by the kidneys over a 6-hour period, resulting in high urinary concentrations. Probenecid slows tubular excretion and produces higher serum levels and increases the duration of measurable serum concentrations.

Cefoxitin passes into pleural and joint fluids and is detectable in antibacterial concentrations in bile. In a published study of geriatric patients ranging in age from 64 to 88 years with normal renal function for their age (creatinine clearance ranging from 31.5 to 174.0 mL/min), the half-life for cefoxitin ranged from 51 to 90 minutes, resulting in higher plasma concentrations than in younger adults. These changes were attributed to decreased renal function associated with the aging process.

Microbiology

Mechanism of Action

Cefoxitin is a bactericidal agent that acts by inhibition of bacterial cell wall synthesis. Cefoxitin has activity in the presence of some beta-lactamases, both penicillinases and cephalosporinases, of Gram-negative and Gram-positive bacteria.

Mechanism of Resistance

Resistance to cefoxitin is primarily through hydrolysis by beta-lactamase, alteration of penicillin-binding proteins (PBPs), and decreased permeability.

Cefoxitin has been shown to be active against most isolates of the following bacteria, both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE**section:

Gram-positive bacteria

Staphylococcus aureus (methicillin-susceptible isolates only)

Staphylococcus epidermidis (methicillin-susceptible isolates only)

Streptococcus agalactiae

Streptococcus pneumoniae

Streptococcus pyogenes

Gram-negative bacteria

Escherichia coli

Haemophilus influenzae

Klebsiella spp.

Morganella morganii

Neisseria gonorrhoeae

Proteus mirabilis

Proteus vulgaris

Providencia spp.

Anaerobic bacteria

Clostridium spp.

Peptococcus niger

Peptostreptococcus spp.

Bacteroides spp.

The following in vitro data are available, but their clinical significance is unknown.

At least 90 percent of the following microorganisms exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for cefoxitin. However, the efficacy of cefoxitin in treating clinical infections due to these microorganisms has not been established in adequate and well-controlled clinical trials.

Gram-negative bacteria

Eikenella corrodens (non-β-lactamase producers)

Anaerobic bacteria

Clostridium perfringens

Prevotella bivia

Susceptibility Test Methods

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: https://www.fda.gov/STIC.

INDICATIONS AND USAGE

Treatment

Cefoxitin for Injection, USP is indicated for the treatment of serious infections caused by susceptible strains of the designated microorganisms in the diseases listed below.

(1) **Lower respiratory tract infections,** including pneumonia and lung abscess, caused by *Streptococcus pneumoniae*, other streptococci (excluding enterococci, e.g., *Enterococcus faecalis* [formerly *Streptococcus faecalis*]), *Staphylococcus aureus* (including penicillinase-producing strains), *Escherichia coli, Klebsiella species*,

Haemophilus influenzae, and Bacteroides species.

- (2) **Urinary tract infections** caused by *Escherichia coli, Klebsiella* species, *Proteus mirabilis, Morganella morganii, Proteus vulgaris* and *Providencia* species (including *P. rettgeri*).
- (3) **Intra-abdominal infections,** including peritonitis and intra-abdominal abscess, caused by *Escherichia coli, Klebsiella* species, *Bacteroides* species including *Bacteroides* fragilis, and *Clostridium* species.
- (4) **Gynecological infections,** including endometritis, pelvic cellulitis, and pelvic inflammatory disease caused by *Escherichia coli, Neisseria gonorrhoeae* (including penicillinase-producing strains), *Bacteroides* species including *B. fragilis, Clostridium* species, *Peptococcus niger, Peptostreptococcus* species and *Streptococcus agalactiae*. Cefoxitin for Injection, USP, like cephalosporins, has no activity against *Chlamydia trachomatis*. Therefore, when Cefoxitin for Injection, USP is used in the treatment of patients with pelvic inflammatory disease and *C. trachomatis* is one of the suspected pathogens, appropriate anti-chlamydial coverage should be added.
- (5) **Septicemia caused by Streptococcus pneumoniae,** *Staphylococcus aureus* (including penicillinase-producing strains), *Escherichia coli, Klebsiella* species, and *Bacteroides* species including *B. fragilis*.
- (6) **Bone and joint infections** caused by *Staphylococcus aureus* (including penicillinase-producing strains).
- (7) **Skin and skin structure infections** caused by *Staphylococcus aureus* (including penicillinase-producing strains), *Staphylococcus epidermidis, Streptococcus pyogenes* and other *streptococci* (excluding enterococci e.g., *Enterococcus faecalis* [formerly *Streptococcus faecalis*]), *Escherichia coli, Proteus mirabilis, Klebsiella* species, *Bacteroides* species including *B. fragilis, Clostridium* species, *Peptococcus niger*, and *Peptostreptococcus* species.

Appropriate culture and susceptibility studies should be performed to determine the susceptibility of the causative organisms to cefoxitin. Therapy may be started while awaiting the results of these studies.

In randomized comparative studies, cefoxitin and cephalothin were comparably safe and effective in the management of infections caused by gram-positive cocci and gram-negative rods susceptible to the cephalosporins. Cefoxitin Injection, USP has a high degree of stability in the presence of bacterial beta-lactamases, both penicillinases and cephalosporinases.

Many infections caused by aerobic and anaerobic gram-negative bacteria resistant to some cephalosporins respond to cefoxitin. Similarly, many infections caused by aerobic and anaerobic bacteria resistant to some penicillin antibiotics (ampicillin, carbenicillin, penicillin G) respond to treatment with Cefoxitin Injection, USP. Many infections caused by mixtures of susceptible aerobic and anaerobic bacteria respond to treatment with Cefoxitin Injection, USP.

Prevention

Cefoxitin for Injection is indicated for the prophylaxis of infection in patients undergoing uncontaminated gastrointestinal surgery, vaginal hysterectomy, abdominal hysterectomy, or cesarean section.

If there are signs of infection, specimens for culture should be obtained for identification of the causative organism so that appropriate treatment may be instituted.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefoxitin for injection and other antibacterial drugs, Cefoxitin for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

Cefoxitin for Injection is contraindicated in patients who have shown hypersensitivity to cefoxitin and the cephalosporin group of antibiotics.

WARNINGS

BEFORE THERAPY WITH CEFOXITIN FOR INJECTION IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEFOXITIN, CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. THIS PRODUCT SHOULD BE GIVEN WITH CAUTION TO PENICILLINSENSITIVE PATIENTS. ANTIBIOTICS SHOULD BE ADMINISTERED WITH CAUTION TO ANY PATIENT WHO HAS DEMONSTRATED SOME FORM OF ALLERGY, PARTICULARLY TO DRUGS. IF AN ALLERGIC REACTION TO CEFOXITIN FOR INJECTION OCCURS, DISCONTINUE THE DRUG. SERIOUS HYPERSENSITIVITY REACTIONS MAY REQUIRE EPINEPHRINE AND OTHER EMERGENCY MEASURES.

Clostridium difficile associated diarrhea (CDAD) has been reported with the use of nearly all antibacterial agents, including Cefoxitin for Injection, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD.

Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

PRECAUTIONS

General

The total daily dose should be reduced when Cefoxitin for Injection is administered to patients with transient or persistent reduction of urinary output due to renal insufficiency (see **DOSAGE AND ADMINISTRATION**), because high and prolonged serum antibiotic concentrations can occur in such individuals from usual doses.

Antibiotics (including cephalosporins) should be prescribed with caution in individuals with a history of gastrointestinal disease, particularly colitis.

As with other antibiotics, prolonged use of Cefoxitin for Injection may result in overgrowth of nonsusceptible organisms. Repeated evaluation of the patient's condition is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Prescribing Cefoxitin for Injection in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for Patients

Patients should be counseled that antibacterial drugs including Cefoxitin for Injection should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Cefoxitin for Injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed.

Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Cefoxitin for Injection or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics, which usually ends when the antibiotic is discontinued. Sometimes after starting the treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Laboratory Tests

As with any potent antibacterial agent, periodic assessment of organ system functions, including renal, hepatic, and hematopoietic, is advisable during prolonged therapy.

Drug Interactions

Increased nephrotoxicity has been reported following concomitant administration of cephalosporins and aminoglycoside antibiotics.

Drug/Laboratory Test Interactions

As with cephalothin, high concentrations of cefoxitin (>100 micrograms/mL) may interfere with measurement of serum and urine creatinine levels by the Jaffé reaction, and produce false increases of modest degree in the levels of creatinine reported. Serum samples from patients treated with cefoxitin should not be analyzed for creatinine if withdrawn within 2 hours of drug administration.

High concentrations of cefoxitin in the urine may interfere with measurement of urinary 17-hydroxycorticosteroids by the Porter-Silber reaction, and produce false increases of modest degree in the levels reported.

A false-positive reaction for glucose in the urine may occur. This has been observed with CLINITEST[†] reagent tablets.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed with cefoxitin to evaluate carcinogenic or mutagenic potential. Studies in rats treated intravenously with 400 mg/kg of cefoxitin (approximately three times the maximum recommended human dose) revealed no effects on fertility or mating ability.

Pregnancy

Reproduction studies performed in rats and mice at parenteral doses of approximately one to seven and one-half times the maximum recommended human dose did not reveal teratogenic or fetal toxic effects, although a slight decrease in fetal weight was observed.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

In the rabbit, cefoxitin was associated with a high incidence of abortion and maternal death. This was not considered to be a teratogenic effect but an expected consequence of the rabbit's unusual sensitivity to antibiotic-induced changes in the population of the microflora of the intestine.

Nursing Mothers

Cefoxitin is excreted in human milk in low concentrations. Caution should be exercised when cefoxitin is administered to a nursing woman.

Pediatric Use

Safety and efficacy in pediatric patients from birth to three months of age have not yet been established. In pediatric patients three months of age and older, higher doses of cefoxitin have been associated with an increased incidence of eosinophilia and elevated SGOT.

Geriatric Use

Of the 1,775 subjects who received cefoxitin in clinical studies, 424 (24%) were 65 and over, while 124 (7%) were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out (see **CLINICAL PHARMACOLOGY**).

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in

dose selection, and it may be useful to monitor renal function (see **DOSAGE AND ADMINISTRATION** and **PRECAUTIONS**).

ADVERSE REACTIONS

Cefoxitin for Injection is generally well tolerated. The most common adverse reactions have been local reactions following intravenous injection. Other adverse reactions have been encountered infrequently.

Local Reactions

Thrombophlebitis has occurred with intravenous administration.

Allergic Reactions

Rash (including exfoliative dermatitis and toxic epidermal necrolysis), urticaria, flushing, pruritus, eosinophilia, fever, dyspnea, and other allergic reactions including anaphylaxis, interstitial nephritis and angioedema have been noted.

Cardiovascular

Hypotension.

Gastrointestinal

Diarrhea, including documented pseudomembranous colitis which can appear during or after antibiotic treatment. Nausea and vomiting have been reported rarely.

Neuromuscular

Possible exacerbation of myasthenia gravis.

Blood

Eosinophilia, leukopenia including granulocytopenia, neutropenia, anemia, including hemolytic anemia, thrombocytopenia, and bone marrow depression. A positive direct Coombs test may develop in some individuals, especially those with azotemia.

Liver Function

Transient elevations in SGOT, SGPT, serum LDH, and serum alkaline phosphatase; and jaundice have been reported.

Renal Function

Elevations in serum creatinine and/or blood urea nitrogen levels have been observed. As with the cephalosporins, acute renal failure has been reported rarely. The role of cefoxitin in changes in renal function tests is difficult to assess, since factors predisposing to prerenal azotemia or to impaired renal function usually have been present.

In addition to the adverse reactions listed above which have been observed in patients treated with cefoxitin, the following adverse reactions and altered laboratory test results have been reported for cephalosporin class antibiotics: Urticaria, erythema multiforme,

Stevens-Johnson syndrome, serum sickness-like reactions, abdominal pain, colitis, renal dysfunction, toxic nephropathy, false-positive test for urinary glucose, hepatic dysfunction including cholestasis, elevated bilirubin, aplastic anemia, hemorrhage, prolonged prothrombin time, pancytopenia, agranulocytosis, superinfection, vaginitis including vaginal candidiasis.

Several cephalosporins have been implicated in triggering seizures, particularly in patients with renal impairment when the dosage was not reduced. (See **DOSAGE AND ADMINISTRATION.**) If seizures associated with drug therapy occur, the drug should be discontinued. Anticonvulsant therapy can be given if clinically indicated.

To report **SUSPECTED ADVERSE REACTIONS**, contact Hikma Pharmaceuticals USA Inc. at 1-877-845-0689 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

The acute intravenous LD_{50} in the adult female mouse and rabbit was about 8.0 g/kg and greater than 1.0 g/kg, respectively. The acute intraperitoneal LD_{50} in the adult rat was greater than 10.0 g/kg.

DOSAGE AND ADMINISTRATION

The intent of this Pharmacy Bulk Package is for the preparation of solutions for intravenous infusion only.

TREATMENT

Adults

The usual adult dosage range is 1 gram to 2 grams every six to eight hours. Dosage should be determined by susceptibility of the causative organisms, severity of infection, and the condition of the patient (see Table 3 for dosage guidelines).

If *C. trachomatis* is a suspected pathogen, appropriate anti-chlamydial coverage should be added, because cefoxitin sodium has no activity against this organism.

Cefoxitin for Injection may be used in patients with reduced renal function with the following dosage adjustments:

In adults with renal insufficiency, an initial loading dose of 1 gram to 2 grams may be given. After a loading dose, the recommendations for maintenance dosage (Table 4) may be used as a guide.

When only the serum creatinine level is available, the following formula (based on sex, weight, and age of the patient) may be used to convert this value into creatinine clearance. The serum creatinine should represent a steady state of renal function.

Males: Weight (kg) x (140 - age)

72 x serum creatinine (mg/100 mL)

Females: 0.85 x above value

In patients undergoing hemodialysis, the loading dose of 1 gram to 2 grams should be given after each hemodialysis, and the maintenance dose should be given as

indicated in Table 4.

Antibiotic therapy for group A beta-hemolytic streptococcal infections should be maintained for at least 10 days to guard against the risk of rheumatic fever or glomerulonephritis. In staphylococcal and other infections involving a collection of pus, surgical drainage should be carried out where indicated.

Pediatric Patients

The recommended dosage in pediatric patients three months of age and older is 80 to 160 mg/kg of body weight per day divided into four to six equal doses. The higher dosages should be used for more severe or serious infections. The total daily dosage should not exceed 12 grams.

At this time no recommendation is made for pediatric patients from birth to three months of age (see **PRECAUTIONS**).

In pediatric patients with renal insufficiency, the dosage and frequency of dosage should be modified consistent with the recommendations for adults (see Table 4).

PREVENTION

Effective prophylactic use depends on the time of administration. Cefoxitin for Injection usually should be given one-half to one hour before the operation, which is sufficient time to achieve effective levels in the wound during the procedure. Prophylactic administration should usually be stopped within 24 hours since continuing administration of any antibiotic increases the possibility of adverse reactions but, in the majority of surgical procedures, does not reduce the incidence of subsequent infection.

For prophylactic use in uncontaminated gastrointestinal surgery, vaginal hysterectomy, or abdominal hysterectomy, the following doses are recommended:

Adults:

2 grams administered intravenously just prior to surgery (approximately one-half to one hour before the initial incision) followed by 2 grams every 6 hours after the first dose for no more than 24 hours.

Pediatric Patients (3 months and older):

30 to 40 mg/kg doses may be given at the times designated above.

Cesarean section patients:

For patients undergoing cesarean section, either a single 2 gram dose administered intravenously as soon as the umbilical cord is clamped OR a 3-dose regimen consisting of 2 grams given intravenously as soon as the umbilical cord is clamped followed by 2 grams 4 and 8 hours after the initial dose is recommended. (See **CLINICAL STUDIES.**)

Table 3 - Guidelines for Dosage of Cefoxitin for Injection

I VDE OT INTECTION	Daily Dosage	Frequency and Route
Uncomplicated forms* of infections such as pneumonia,	3-4	1 gram every 6-8

urinary tract infection, cutaneous infection	grams	hours IV
Moderately severe or severe infections	6-8 grams	1 gram every 4 hours or 2 grams every 6-8 hours IV
Infections commonly needing antibiotics in higher dosage (e.g., gas gangrene)	12 grams	2 grams every 4 hours or 3 grams every 6 hours IV

^{*}Including patients in whom bacteremia is absent or unlikely.

Table 4 - Maintenance Dosage of Cefoxitin for Injection in Adults with Reduced Renal Function

Kenal Function	Creatinine Clearance (mL/min)	Dose (Grams)	Frequency
Mild impairment Moderate impairment Severe impairment Essentially no function	50 - 30 29 - 10 9 - 5 < 5	1 - 2 1 - 2 0.5 - 1 0.5 - 1	Every 8-12 hours Every 12- 24 hours Every 12-24 hours Every 24-48 hours

Table 5 - Preparation of Solution for Intravenous Administration

	Added		Approximate Average concentration (mg/mL)
10 grams Pharmacy Bulk Package	43 or 93	49 or 98.5	200 or 100

^{*}Shake to dissolve and let stand until clear

DIRECTIONS FOR PROPER USE OF PHARMACY BULK PACKAGE BOTTLE - NOT FOR DIRECT INFUSION

For Bulk Packages

The Pharmacy Bulk Package bottle is for use in a pharmacy admixture service under a laminar flow hood. Penetration into the Pharmacy Bulk Package bottle should be made

only one time after reconstitution with a sterile transfer set or other sterile dispensing device, which allows measured distribution of the contents. Dispense the contents in aliquots using aseptic technique. The use of a syringe with a needle is not recommended as it may cause leakage (see **DOSAGE AND ADMINISTRATION**).

AFTER INITIAL ENTRY USE ENTIRE CONTENTS OF THE PHARMACY BULK PACKAGE PROMPTLY. A maximum time of **4 HOURS** from initial entry is permitted to complete fluid transfer operations. ANY UNUSED PORTION MUST BE DISCARDED WITHIN **4 HOURS.** This time limit should begin with the introducing of solvent or diluent into the Pharmacy Bulk Package bottle. **RECONSTITUTED BULK SOLUTION SHOULD NOT BE USED FOR DIRECT INFUSION.**

PREPARATION OF SOLUTION

Table 5 is provided for convenience in constituting Cefoxitin for Injection Pharmacy Bulk Package.

The 10 grams bulk packages should be constituted with 43 mL or 93 mL of Sterile Water for Injection, Bacteriostatic Water for Injection, 0.9 percent Sodium Chloride Injection, or 5 percent Dextrose Injection. **RECONSTITUTED STOCK SOLUTION MUST BE TRANSFERRED AND FURTHER DILUTED FOR I.V. INFUSION.** These primary solutions may be further diluted in 50 to 1000 mL of the diluents listed under the **COMPATIBILITY AND STABILITY** section.

Benzyl alcohol as a preservative has been associated with toxicity in neonates. While toxicity has not been demonstrated in pediatric patients greater than three months of age, in whom use of Cefoxitin for Injection may be indicated, small pediatric patients in this age range may also be at risk for benzyl alcohol toxicity. Therefore, diluent containing benzyl alcohol should not be used when Cefoxitin for Injection is constituted for administration to pediatric patients in this age range.

ADMINISTRATION

Cefoxitin for Injection may be administered intravenously after constitution.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

The intent of this pharmacy bulk package is for the preparation of solutions for intravenous infusion only.

Intravenous Administration

The intravenous route is preferable for patients with bacteremia, bacterial septicemia, or other severe or life-threatening infections, or for patients who may be poor risks because of lowered resistance resulting from such debilitating conditions as malnutrition, trauma, surgery, diabetes, heart failure, or malignancy, particularly if shock is present or impending.

For intermittent intravenous administration, a solution containing 1 gram or 2 grams in 10 mL of Sterile Water for Injection can be injected over a period of three to five minutes. Using an infusion system, it may also be given over a longer period of time through the tubing system by which the patient may be receiving other intravenous solutions. However, during infusion of the solution containing Cefoxitin for Injection, it is advisable to temporarily discontinue administration of any other solutions at the same

site.

For the administration of higher doses by continuous intravenous infusion, a solution of Cefoxitin for Injection may be added to an intravenous bottle containing 5 percent Dextrose Injection, 0.9 percent Sodium Chloride Injection, or 5 percent Dextrose and 0.9 percent Sodium Chloride Injection. BUTTERFLY^{®††} or scalp vein-type needles are preferred for this type of infusion.

Solutions of Cefoxitin for Injection, like those of most beta-lactam antibiotics, should not be added to aminoglycoside solutions (e.g., gentamicin sulfate, tobramycin sulfate, amikacin sulfate) because of potential interaction. However, cefoxitin for injection and aminoglycosides may be administered separately to the same patient.

COMPATIBILITY AND STABILITY

Cefoxitin for Injection supplied in pharmacy bulk package bottles and constituted to 1 gram/10 mL with Sterile Water for Injection, Bacteriostatic Water for Injection, (see **PREPARATION OF SOLUTION**), 0.9 percent Sodium Chloride Injection, or 5 percent Dextrose Injection, should be DISCARDED 4 HOURS AFTER INITIAL ENTRY. FURTHER DILUTION IS REQUIRED BEFORE USE. **RECONSTITUTED BULK SOLUTION SHOULD NOT BE USED FOR DIRECT INFUSION.**

These primary solutions may be further diluted in 50 mL to 1000 mL of the following diluents and maintain potency for an additional 18 hours at room temperature or an additional 48 hours under refrigeration:

0.9 percent Sodium Chloride Injection

5 percent or 10 percent Dextrose Injection

5 percent Dextrose and 0.9 percent Sodium Chloride Injection

5 percent Dextrose Injection with 0.2 percent or 0.45 percent saline solution

Lactated Ringer's Injection

5 percent Dextrose in Lactated Ringer's Injection

5 percent Sodium Bicarbonate Injection

M/6 sodium lactate solution

Mannitol 5% and 10%

After the periods mentioned above, any unused solutions should be discarded.

HOW SUPPLIED

Cefoxitin for Injection, USP is a dry white to off-white powder supplied in Pharmacy Bulk Package bottle. Each Pharmacy Bulk Package bottle contains cefoxitin sodium as follows:

Cefoxitin for Injection, USP 10 g, Pharmacy Bulk Package - (10 gram cefoxitin equivalent)

NDC 0143-9876-10 cartons of 10 pharmacy bulk packages

Special storage instructions

Cefoxitin for Injection in the dry state should be stored between 2º to 25°C (36º to 77°F). Avoid exposure to temperatures above 50°C. The dry material as well as solutions tend to darken, depending on storage conditions; product potency, however, is not adversely affected.

CLINICAL STUDIES

A prospective, randomized, double-blind, placebo-controlled clinical trial was conducted to determine the efficacy of short-term prophylaxis with cefoxitin in patients undergoing cesarean section who were at high risk for subsequent endometritis because of ruptured membranes. Patients were randomized to receive either three doses of placebo (n=58), a single dose of cefoxitin (2 g) followed by two doses of placebo (n=64), or a three-dose regimen of cefoxitin (each dose consisting of 2 g) (n=60), given intravenously, usually beginning at the time of clamping of the umbilical cord, with the second and third doses given 4 and 8 hours post-operatively. Endometritis occurred in 16/58 (27.6%) patients given placebo, 5/63 (7.9%) patients given a single dose of cefoxitin, and 3/58 (5.2%) patients given three doses of cefoxitin. The differences between the two groups treated with cefoxitin and placebo with respect to endometritis were statistically significant (p<0.01) in favor of cefoxitin. The differences between the one-dose and three-dose regimens of cefoxitin were not statistically significant.

Two double-blind, randomized studies compared the efficacy of a single 2 gram intravenous dose of cefoxitin to a single 2 gram intravenous dose of cefotetan in the prevention of surgical site-related infection (major morbidity) and non-site-related infections (minor morbidity) in patients following cesarean section. In the first study, 82/98 (83.7%) patients treated with cefoxitin and 71/95 (74.7%) patients treated with cefotetan experienced no major or minor morbidity. The difference in the outcomes in this study (95% CI: -0.03, +0.21) was not statistically significant. In the second study, 65/75 (86.7%) patients treated with cefoxitin and 62/76 (81.6%) patients treated with cefotetan experienced no major or minor morbidity. The difference in the outcomes in this study (95% CI: -0.08, +0.18) was not statistically significant.

In clinical trials of patients with intra-abdominal infections due to *Bacteroides fragilis* group microorganisms, eradication rates at 1 to 2 weeks post treatment for isolates were in the range of 70% to 80%. Eradication rates for individual species are listed below:

Bacteroides distasonis	7/10	(70%)
Bacteroides fragilis	26/33	(79%)
Bacteroides ovatus	10/13	(77%)
B. thetaiotaomicron	13/18	(72%)

REFERENCES

1. Carver PL, Nightingale CH and Quintiliani R. Pharmacokinetics and pharmacodynamics of total and unbound cefoxitin and cefotetan in healthy volunteers. Journal of Antimicrobial Chemotherapy (1989) 23, 99-106.

Manufactured by:

HIKMA FARMACÊUTICA (PORTUGAL), S.A.

Estrada do Rio da Mó, nº 8, 8A e 8B - Fervença,

2705 - 906 Terrugem SNT

PORTUGAL

Distributed by:

Hikma Pharmaceuticals USA Inc.

Berkeley Heights, NJ 07922

Revised June 2020

PIN146-WES/6

PRINCIPAL DISPLAY PANEL

NDC 0143-**9876**-01 Rx only

CEFOXITIN

FOR INJECTION, USP

PHARMACY BULK PACKAGE NOT FOR DIRECT INFUSION

10 grams* per Pharmacy Bulk Package

Bottle contains cefoxitin sodium powder equivalent to 10 g cefoxitin

For Intravenous use ONLY



CEFOXITIN FOR INJECTION, USP PHARMACY BULK PACKAGE NOT FOR DIRECT INFUSION 10 grams per Pharmacy Bulk Package For Intravenous use ONLY



CEFOXITIN

cefoxitin injection, powder, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9876
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CEFOXITIN SODIUM (UNII: Q68050H03T) (CEFOXITIN - UNII:60EV9DX57Y)	CEFOXITIN	10 g

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0143-9876- 10	10 in 1 CARTON	02/26/2010		
1	NDC:0143-9876- 01	1 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065239	02/26/2010	

Labeler - Hikma Pharmaceuticals USA Inc. (001230762)

Establishment				
Na me	Address	ID/FEI	Business Operations	
HIKMA FARMACEUTICA (PORTUGAL), S.A		452742943	manufacture(0143-9876)	

Revised: 3/2025 Hikma Pharmaceuticals USA Inc.