

NOHIST DM- chlorpheniramine maleate, dextromethorphan hydrobromide and phenylephrine hydrochloride liquid
Larken Laboratories, Inc.

NoHist DM

Drug Facts

Active Ingredients

(In each 5 mL teaspoonful)

Chlorpheniramine Maleate, USP 4 mg

Dextromethorphan HBr, USP 15 mg

Phenylephrine HCl, USP 10 mg

Purpose

Chlorpheniramine Maleate Antihistamine

Dextromethorphan HBr Antitussive (cough suppressant)

Phenylephrine HCl Nasal decongestant

Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis):

- cough due to minor throat and bronchial irritation
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking any other nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product**Do not exceed recommended dosage.**

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur.
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Keep out of the reach of children

In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

Do not exceed 6 doses in a 24-hour period

Age	Dose
Adults and children over 12	1 teaspoonsful (5 mL) every 4

years of age	hours
Children 6 to under 12 years of age	1/2 teaspoonsful (2.5 mL) every 4 hours
Children under 6 years of age	Ask your doctor

Other Information

- store at 20°-25°C (68°-77°F)
- very low sodium, contains 5 mg sodium per 5 mL teaspoonful

Inactive Ingredients

citric acid, edetate disodium, glycerin, grape flavoring, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, and sodium citrate dihydrate

Questions or Comments

Call 1-601-855-7678 weekdays from 9:00 am to 4:00 pm CST or go to <http://www.larkenlabs.com>.

Principal Display Panel

Figure 1: 16 oz. Bottle Label

NDC 68047-186-16

NOHIST•DM

ANTI-HISTAMINE / ANTITUSSIVE
NASAL DECONGESTANT

SUGAR FREE / ALCOHOL FREE
DYE FREE

Grape Flavored Liquid

**DO NOT USE IF FOIL SEAL UNDER
THE CAP IS BROKEN OR MISSING.**

Distributed by:

**LARKEN
LABORATORIES**

Canton, MS 39046

16 fl. oz. (473 mL)



Lot/Exp. date:

Drug Facts

Active ingredients (in each 5 mL teaspoonful)	Purpose
Chlorpheniramine Maleate, USP 4 mg	Antihistamine
Dextromethorphan HBr, USP 15 mg	Antitussive (cough suppressant)
Phenylephrine HCl, USP 10 mg	Nasal decongestant

Uses
temporarily relieves these symptoms due to hay fever (allergic rhinitis): ■ cough due to minor throat and bronchial irritation ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat
■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ temporarily restores freer breathing through the nose

Warnings
Do not use
■ to sedate a child or make a child sleepy
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

Ask a doctor before use if you have
■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema ■ a cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
■ taking any other nasal decongestant or stimulant
■ taking sedatives or tranquilizers

When using this product
Do not exceed recommended dosage.
■ marked drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children.

200964-01 Rev. 09/2025

PEEL

NOHIST DM

chlorpheniramine maleate, dextromethorphan hydrobromide and phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68047-186
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE -	CHLORPHENIRAMINE	4 mg

UNII:3U6IO1965U)	MALEATE	in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68047-186-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/06/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/06/2011	

Labeler - Larken Laboratories, Inc. (149484540)

Registrant - Larken Laboratories, Inc. (149484540)