

**PETARMOR SURE SHOT 2X- pyrantel pamoate liquid**  
**Sergeant's Pet Care Products, LLC**

-----  
**PETARMOR®**  
**FOR DOGS**  
**Sure Shot® 2X**  
**(pyrantel pamoate)**

**Extended Content Label Panel FR3**

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly internals to prevent *T. canis* reinfestation.

**RECOMMENDED STORAGE: STORE BELOW 86°F (30°C).**

**PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER**

**petarmor.com | @petarmor**

**PETARMOR FOR DOGS**

NDC 21091-716-01

**Sure Shot® 2X**

*(pyrantel pamoate oral suspension)*

**Canine Anthelmintic Suspension**

Active Ingredient: 4.54 mg of Pyrantel Base as  
Pyrantel Pamoate per mL

**NOT FOR USE IN HUMANS**

Approved by FDA under ANADA # 200-007

**NET CONTENTS - 2 fl. oz. (60 mL)**

**KEEP OUT OF REACH OF CHILDREN. Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.**

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory

fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

Read accompanying professional information under this label.

**Manufactured for:**

**Sergeant's Pet Care Products, LLC**  
**Omaha, NE 68138**

P01063V02 01/2025

Take Time (Eye Symbol) Observe Label Directions

**READ PACKAGE EXTENDED CONTENT LABEL FOR FULL PRESCRIBING INFORMATION**

(PEEL)

**Extended Content Label Panel BK3**


**Sure Shot® 2X** (pyrantel pamoate oral suspension)

**Canine Anthelmintic Suspension**

**NOT FOR USE IN HUMANS**

**DESCRIPTION:** Sure Shot® 2X Liquid Wormer is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given to the right:



**(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4,4' methylenebis [3-hydroxy-2-naphthoate](1:1)**

**INDICATIONS AND USAGE:** Sure Shot® 2X suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

## Extended Content Label Panel FR2

Sure Shot<sup>®</sup> 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**PRECAUTIONS:** THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, **IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.**

### KEEP OUT OF REACH OF CHILDREN

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of bodyweight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**SAFETY AND EFFICACY STUDIES:** Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD(50) of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 138 times

## Extended Content Label Panel BK2

the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud.

Additional data has demonstrated the safe use of pyrantel pamoate in dogs having heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C). Keep Sure Shot<sup>®</sup> 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

HOW SUPPLIED: Sure Shot® 2X is supplied in 2 fl. oz. (60 mL) bottle.

Approved by FDA under ANADA # 200-007

Take Time (Eye Symbol) Observe Label Directions

**Contact Information:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Sergeant's Pet Care Products, LLC at 1-800-224-7387. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

### **Extended Content Label Panel FR1 (Bottom Ply)**

#### **IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly internals to prevent *T. canis* reinfestation.

#### **PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER**

PETARMOR®

NDC 21091-716-01

Sure Shot® 2X

*(pyrantel pamoate oral suspension)*

#### **Canine Anthelmintic Suspension**

Active Ingredient: 4.54 mg of Pyrantel Base as

Pyrantel Pamoate per mL

#### **NOT FOR USE IN HUMANS**

Approved by FDA under ANADA # 200-007

**NET CONTENTS - 2 fl. oz. (60 mL)**

**RECOMMENDED STORAGE: STORE BELOW 86°F (30°C).**

**KEEP OUT OF REACH OF CHILDREN. Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.**

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and

control of parasitism. Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping. Read package extended content label for full prescribing Information.

**Manufactured for:**

**Sergeant's Pet Care Products, LLC  
Omaha, NE 68138**

P01063V02 01/2025

Take Time (Eye Symbol) Observe Label Directions

**Primary Label (ECL - Extended Content) - 60 mL Bottle**

FR3

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**RECOMMENDED STORAGE:** STORE BELOW 88°F (30°C).

PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER ▼  
petarmor.com | @petarmor

**PETARMOR.**  
FOR DOGS

NDC 21091-716-01

**Sure Shot® 2X**  
(pyrantel pamoate oral suspension)

**Canine Anthelmintic Suspension**

Active Ingredient: 4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL  
**NOT FOR USE IN HUMANS**  
Approved by FDA under ANADA# 200-007

**NET CONTENTS - 2 fl. oz. (60 mL)**

**KEEP OUT OF REACH OF CHILDREN.** Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping. Read accompanying professional information under this label.

**Manufactured for:**  
Sergeant's Pet Care Products, LLC  
Omaha, NE 68138  
P01063V02 01/2025

Take Time (Eye Symbol) Observe Label Directions

**READ PACKAGE EXTENDED CONTENT LABEL FOR FULL PRESCRIBING INFORMATION**

BK3

**Sure Shot® 2X** (pyrantel pamoate oral suspension)  
Canine Anthelmintic Suspension

**NOT FOR USE IN HUMANS**

**DESCRIPTION:** Sure Shot® 2X Liquid Wormer is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given to the right.

**INDICATIONS AND USAGE:** Sure Shot® 2X suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl]pyrimidine 4,4'-methylenebis[3-hydroxy-2-naphthoate] (1:1)

hinge

FR2

**PROFESSIONAL INFORMATION CONTINUED**

Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**PRECAUTIONS:** THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.

**KEEP OUT OF REACH OF CHILDREN**

**DOSE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**SAFETY AND EFFICACY STUDIES:** Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (95%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD<sub>50</sub> of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 130 times.

hinge

Bk2

the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 54 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud.

Additional data has demonstrated the safe use of pyrantel pamoate in dogs having heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming, such as immunization and antibacterial treatment.

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C). Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

**HOW SUPPLIED:** Sure Shot® 2X is supplied in 2 fl. oz. (60 mL) bottle.

Approved by FDA under ANADA # 200-007

**Take Time**  **Observe Label Directions**

**Contact Information:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Sergeant's Pet Care Products, LLC at 1-800-224-7387. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

hinge

FR1

**IMPORTANT: SHAKE WELL BEFORE USING**

**DOSAGE AND ADMINISTRATION:** Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

**PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER ▼**

**PETARMOR**  
FOR DOGS

NDC 21091-716-01

**Sure Shot® 2X**  
(pyrantel pamoate oral suspension)

**Canine Anthelmintic Suspension**

Active Ingredient: 454 mg of Pyrantel Base as Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS  
Approved by FDA under ANADA # 200-007

**NET CONTENTS - 2 fl. oz. (60 mL)**

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C).

**KEEP OUT OF REACH OF CHILDREN. Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.**

**INDICATIONS AND USAGE:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

Read package extended content label for full prescribing information.

**Manufactured for:**  
Sergeant's Pet Care Products, LLC  
Omaha, NE 68138  
P01063V02 01/2025

**Take Time**  **Observe Label Directions**

hinge

## Secondary Label - Carton



**Sure Shot® 2X**  
(pyrantel pamoate oral suspension)  
Canine Anthelmintic Suspension

**DOSE AND ADMINISTRATION:**  
Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

IMPORTANT: SHAKE WELL BEFORE USE



Read package extended content label for full prescribing information.

petarmor.com | @petarmor

**Sure Shot® 2X**  
(pyrantel pamoate oral suspension)

**INDICATIONS AND USAGE:**  
For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Sure Shot® 2X suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.



P00989 12/2024

**RECOMMENDED STORAGE:** STORE BELOW 86°F (30°C)  
Keep Sure Shot® 2X in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CARTON ▼

**PETARMOR.**  
FOR DOGS

Each 2oz. bottle treats  
**120 lbs.**

**Sure Shot® 2X**  
(pyrantel pamoate oral suspension)



**Canine Anthelmintic Suspension**

Removes Large Roundworms and Hookworms in Puppies & Dogs

**Active Ingredient:**  
4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL  
**NOT FOR USE IN HUMANS**  
**KEEP OUT OF REACH OF CHILDREN**  
Approved by FDA under ANADA # 200-037

NET CONTENTS - 2 fl. oz. (60mL)

**PETARMOR.**  
FOR DOGS

**Sure Shot® 2X**

Distributed by: Sergeant's Pet Care Products, LLC  
10077 S. 134th Street, Omaha, NE 68138  
1-800-224-7387

**PETARMOR SURE SHOT 2X**  
pyrantel pamoate liquid

## Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:21091-716
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PYRANTEL PAMOATE</b> (UNII: 81BK194Z5M) (Pyrantel - UNII:4QIH0N49E7)	Pyrantel	4.54 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>Methylparaben Sodium</b> (UNII: CR6K9C2NHK)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUTTERSCOTCH	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:21091-716-01	1 in 1 CARTON		
1		60 mL in 1 BOTTLE		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200007	10/30/1996	

**Labeler** - Sergeant's Pet Care Products, LLC (876995171)

**Registrant** - Happy Jack (003189354)

## Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
ABS Corporation		048022727	MANUFACTURE, LABEL, PACK

## Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Cosma Spa		428655732	api manufacture

Revised: 8/2025

Sergeant's Pet Care Products, LLC