

COUNTERACT ALLERGY- loratadine tablet
Melaleuca, Inc.

CounterAct Allergy Content of Label

Active ingredient (in each tablet)

Loratadine 10 mg

Purpose

Antihistamine

Use temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- protect from excessive moisture

Inactive ingredients lactose monohydrate, magnesium stearate, povidone, pregelatinized starch

Questions or comments? 1-800-282-3000

Non-Drowsy 24-Hour*
CounterAct Allergy
Loratadine Tablets, 10 mg / Antihistamine
Indoor & Outdoor Allergies



Sneezing
Itchy, Watery Eyes
Runny Nose
Itchy Throat or Nose

**When taken as directed. See Drug Facts panel.*

Purpose
Loratadine 10 mg,Antihistamine
Use temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose. **When using this product** do not take more than directed. Taking more than directed may cause drowsiness. **Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ adults and children 6 years and over: 1 tablet daily; not more than 1 tablet in 24 hours ■ children under 6 years of age: ask a doctor ■ consumers with liver or kidney disease: ask a doctor

Other information ■ do not use if printed foil under cap is broken or missing ■ store at 20° - 25°C (68° - 77°F)


Questions? 208-534-3000

Distrib: Medeva, Inc
Methuen, MA 01842
To order: 208-534-3000
Medeva.com № 654 06/25 U

NOT FOR RESALE



: 61265 MB F1



COUNTERACT ALLERGY

loratadine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54473-166
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII: 7AJ03BO7QN)	LORATADINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (White colored)	Score	no score
Shape	OVAL (Oval shaped)	Size	8mm
Flavor		Imprint Code	L612
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54473-166-32	1 in 1 BOX	02/28/2025	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075209	08/19/2023	

Labeler - Melaleuca, Inc. (139760102)

Revised: 3/2025

Melaleuca, Inc.