## TOPCARE CHILDRENS ALLERGY- diphenhydramine hydrochloride solution Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## Topco Associates LLC. Children's Allergy Liquid Drug Facts

## Active ingredient (in each 5 mL)

Diphenhydramine HCl 12.5 mg

## **Purpose**

**Antihistamine** 

#### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- itching of the nose or throat
- runny nose
- itchy, watery eyes

# Warnings

#### Do not use

- with any other product containing diphenhydramine, even one used on skin
- to make a child sleepy

#### Ask a doctor before use if the child has

- a breathing problem such as chronic bronchitis
- glaucoma
- a sodium-restricted diet

# Ask a doctor or pharmacist before use if the child is

taking sedatives or tranquilizers

# When using this product

- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- find right dose on chart below
- mL = milliliter
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL to 10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F). Protect from light. Store in outer carton until contents used.
- · do not use if printed neckband is broken or missing

# **Inactive ingredients**

anhydrous citric acid, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, poloxamer 407, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution

## Questions or comments?

1-888-423-0139

# **Principal Display Panel**

TopCare® health

COMPARE TO CHILDREN'S BENADRYL® ALLERGY LIQUID ACTIVE INGREDIENT children's Allergy Liquid

**ANTIHISTAMINE** 

## DIPHENHYDRAMINE HCI

# 12.5 mg/5 mL ORAL SOLUTION

# 4-6 HOURS/DOSE

## **RELIEF OF:**

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Ithy Throat or Nose

Alcohol Free

Ages 6 to 11 Years

**CHERRY FLAVOR** 

8 FL OZ (237 mL)



DIPHENHYDRAMINE HCI 12.5 mg/5 mL ORAL SOLUTION



**Alcohol Free** 

- Itchy, Watery EyesItchy Throat or Nose

CHERRY



8 FL OZ (237 mL)



+TopCare.

# children's **Allergy**

**ANTIHISTAMINE** 

DIPHENHYDRAMINE HCI 12.5 mg/5 mL ORAL SOLUTION



RELIEF OF: • Runny Nose • Sneezing

Itchy, Watery EyesItchy Throat or Nose



8 FL OZ (237 mL)

Drug Facts

Active ingredient (in each 5 mL) )iphenhydramine HCl 12.5 mg.

NOT WIS MILE OF THE SOLUTION DICHEMINDS AN INE HO

**Diupi VD19IIA** 

USES temporarity relieves these symptoms due to hay tever or other upper respiratory ellergies:

sneezing the incse or throat

**◆TopCare** 

■ runny nose ■ Itchy, watery eyes

Warnings
Do not use
■ with any other product containing diphenhydramine, even
one used on skin
■ to make a child sleepy

Ask a doctor before use if the child has ■ a breathing problem such as chronic bronchitis

■ glaucoma ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers

When using this product
■ marked drowsiness may occur
■ excitability may occur, especially in children
■ excitability may occur, especially in children
■ sedatives and tranquilizers may increase drowsiness

Keep out of reach of children. In case of overdose, get medical help or contact a Polson Control Center right away (1-800-222-1222).

Directions
■ find right dose on chart below
■ mL = milliliter

take every 4 to 6 hours, or as directed by a doctor do not take more than 6 doses in 24 hours

Age (yr) children under 2 years do not use children 2 to 5 years do not use unless directed by a doctor

children 6 to 11 years 5 mL to 10 mL
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

■ each 5 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F). Protect from light. Store in oute

Inactive ingredients
antivorous citric acid, D&C red #33, FD&C red #40, flavor, ampurous cinic aau, bac fed #35, Foot fed #40, iavor, glycerin, high fructose corn syrup, poloxamer 407, purifier water, sodium benzoate, sodium chioride, sodium citrate, sorbitol solution Drug Facts (continued)

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC ELK GROVE VILLAGE, IL 60007 ©TOPCO PERA0521 QUESTIONS? 1-888-423-0139 topcare@topco.com www.topcarebrand.com



Scan here for more information or call 1-888-423-0139



GLUTEN FREE



This Top Care® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Benadryl® Allergy Liquid.



LOT NO.

EXP.

37934 88 (9

# TOPCARE CHILDRENS ALLERGY

diphenhydramine hydrochloride solution

#### **Product Information Product Type HUMAN OTC DRUG**

**ORAL** 

**Route of Administration** 

Item Code (Source)

NDC:36800-379

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
D&C RED NO. 33 (UNII: 9DBA0SBB0L)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
POLOXAMER 407 (UNII: TUF2IVW3M2)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CHLORIDE (UNII: 451W47IQ8X)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
GLYCERIN (UNII: PDC6A3C0OX)		

Product Characteristics			
Color	RED (Bluish-Red)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-379- 26	1 in 1 CARTON	12/15/1987	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-379- 34	1 in 1 CARTON	11/18/2003	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/15/1987	

Revised: 2/2022 Topco Associates LLC