

EQUALINE EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent
United Natural Foods, Inc. dba UNFI

Equaline Effervescent Cold Relief

Active ingredients

(in each effervescent tablet)

Aspirin 325 mg (NSAID)*

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Antihistamine

Nasal decongestant

Uses

temporarily relieves these symptoms of the common cold:

■ fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache ■ minor aches and pains

■ runny nose ■ sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic. ■ you are taking sedatives or tranquilizers.

You have ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away. ■ pain or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new

symptoms occur ■ redness or swelling is present

■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not use more than directed (see overdose warning)

■ adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours

■ do not take more than 8 tablets in 24 hours

■ *children under 12 years: ask a doctor*

Other information

each tablet contains: sodium 464 mg

■ phenylketonurics: contains phenylalanine 9 mg per tablet

■ store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments?

1-877-932-7948

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription nonsteroidal anti-inflammatory drug (NSAID) (certain drugs for depression, psychiatric medication, osteoarthritis, or Parkinson's disease), or for 2 weeks after the NSAID drug, if you do not know if your prescription drug contains an NSAID, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you are taking sedatives or tranquilizers. You have
- glaucoma
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are:

- presently taking a prescription drug
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools

Drug Facts

Active Ingredients

(In each effervescent tablet)

Purpose

Aspirin 325 mg (NSAID) Pain reliever/fever reducer
 Chlorpheniramine maleate 2 mg Antihistamine
 Phenylephrine bitartrate 7.8 mg Nasal decongestant
 Nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms of the common cold:

- fever
- sneezing
- nasal congestion
- sore throat
- headache
- minor aches and pains
- runny nose

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach bleeding

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed

compare to Alka-Seltzer Plus® Cold active ingredients**

EQUALINE®

original flavor

effervescent cold relief

fast relief of:

- nasal congestion
- runny nose
- sneezing
- headache
- sinus pain & pressure

20 effervescent tablets

NDC 41163-798-20

compare to Alka-Seltzer Plus® Cold active ingredients**

EQUALINE®

original flavor

effervescent cold relief

aspirin (NSAID)
 (pain reliever/fever reducer)
 chlorpheniramine maleate (antihistamine)
 phenylephrine bitartrate (nasal decongestant)

fast relief of:

- nasal congestion
- runny nose
- sneezing
- headache
- sinus pain & pressure

20 effervescent tablets

Drug Facts (continued)

- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
- do not take more than 8 tablets in 24 hours

Drug Facts (continued)

- children under 12 years: ask a doctor

Other information

each tablet contains:

- sodium 464 mg
- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments?


1-877-932-7948

DISTRIBUTED BY SUPERVALU INC.
 EDEN PRAIRIE, MN 55344 USA
 MADE IN USA
 Contact us at 1-877-932-7948, or
 www.supervalu-ourownbrands.com
 GLUTEN FREE
 **This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold.

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

PN: 87360 EL C4
 REV: 08/19

0 41163 40547 9



EQUALINE EFFERVESCENT COLD RELIEF

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-768
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
MANNITOL (UNII: 3OWL53L36A)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	CF
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-768-20	10 in 1 CARTON	09/24/2019	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M012	05/29/2014	
--------------------	------	------------	--

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 11/2024

United Natural Foods, Inc. dba UNFI