

LOPERAMIDE HYDROCHLORIDE - loperamide hydrochloride tablet
Amerisource Bergen

Drug Facts

Active ingredient (in each tablet)

Loperamide hydrochloride USP 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 tablets after the first loose stool; 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9 to 11 years (60 to 95 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours
children 6 to 8 years (48 to 59 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children 2 to 5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20° to 25°C (68° to 77°F).
- **do not use if carton or blister unit is open or torn**
- Meets USP dissolution test 2
- see side panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, D & C yellow No. 10 aluminum lake, FD & C blue No. 1, lactose

monohydrate, magnesium stearate, microcrystalline cellulose, and sodium starch glycolate

Questions or comments?

call **1-855-274-4122**

Distributed By
AmerisourceBergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?

www.mygnp.com

Made in India

Code : TS/DRUGS/22/2009

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 2 mg Blister Carton 12(2 x 6's Tablets)

**GOOD
NEIGHBOR
PHARMACY®**

***Compare to Imodium® A-D
active ingredient
NDC 46122-738-53**

**Loperamide Hydrochloride
Tablets USP 2 mg**

Anti-Diarrheal

Controls the symptoms of diarrhea

12 Tablets

actual size



ABCF-10276227



Compare to Imodium® A-D active ingredient

NDC 46122-738-53

Loperamide Hydrochloride
Tablets USP 2 mg

Anti-Diarrheal

Controls the symptoms of diarrhea

12 Tablets

actual size



This product is not manufactured or distributed by
Pfizer Consumer Healthcare, owner of the registered
trademark Imodium® A-D.

Lot: P1050530

LM-5068

NVZ

Batch Coding Area

Distributed By
AmisourceBerg
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

Made in India
Code: TS/D/PL/GS/22/2008

Drug Facts (continued)

Active ingredient (in each tablet): Purpose
Loperamide hydrochloride USP 2 mg.....Anti-diarrheal

Warnings
Allergy alert: Do not use if you have ever had rash or other allergic reaction to loperamide hydrochloride


Use
controls symptoms of diarrhea, including travelers' diarrhea

Do not use if you have blood or black stool
liver disease ■ a history of abnormal heart rhythm
Ask a doctor or pharmacist before use if you are taking a prescription drug.
Loperamide may interact with certain prescription drugs.
When using this product, dizziness, or weakness or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.
If pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions
■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ The right dose on chart. If possible, use weight to dose; otherwise, use age.
adults and children 12 years and over
1 tablet after the first loose stool; 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9 to 11 years (60 to 95 lbs)
1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours
children 6 to 8 years (48 to 59 lbs)
1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children 2 to 5 years (34 to 47 lbs)
ask a doctor
children under 2 years (up to 33 lbs)
do not use

Other information
■ Store at 20° to 25° C (68° to 77° F). ■ Do not use if carton or blister unit is open or damaged.
■ Meets USP Dissolution Test 2 ■ See side panel for lot number and expiration date



LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-738
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	GREEN (Light Green)	Score	2 pieces
Shape	CAPSULE (Biconvex)	Size	10mm
Flavor		Imprint Code	L;28
Contains			

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-738-53	2 in 1 CARTON	07/05/2023	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:46122-738-62	4 in 1 CARTON	07/20/2023	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206548	07/05/2023	

Labeler - Amerisource Bergen (007914906)

Registrant - Aurohealth LLC (078728447)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(46122-738) , MANUFACTURE(46122-738)

Revised: 3/2024

Amerisource Bergen