

**GERI-TUSSIN DM- dextromethorphan hydrobromide and guaifenesin liquid**  
**Geri-Care Pharmaceuticals, Corp**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**GERI-TUSSIN DM**

**Active Ingredients (per teaspoonful=5 mL)**

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

**Purpose**

Expectorant

Cough Suppressant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Do not use** if you are hypersensitive to any of the ingredients.

**Ask a doctor before use if you have**

- a cough with too much phlegm (mucus) that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- a chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or accompanied by fever, rash, or headaches that lasts. These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take every 4 hours as needed, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- do not exceed recommended dose

Adult and children 12 years and over

2 teaspoonfuls (10 mL)

Children under 12 years	consult a doctor
-------------------------	------------------

**Inactive ingredients**

citric acid, FD&C Red #40, flavors, glucose, glycerin, high fructose corn syrup, menthol, saccharin sodium, sodium benzoate, water

**Other information**

- store at room temperature between 15°-30°C (59° -86°F)
- protect from freezing.
- protect from light.
- Each teaspoon (5 mL) contains: sodium 2 mg.
- for institutional use only

TAMPER EVIDENT: Do not use this product if breakaway band on cap is broken or missing

**Distributed by:**

GERI-CARE PHARMACEUTICALS CORP.

1650 63rd Street

Brooklyn, NY 11204

# GERI-CARE

NDC 57896-660-16

## GERI-TUSSIN DM

**COUGH SUPPRESSANT • EXPECTORANT**

• **CONTROLS COUGH**

• **LOOSENS & RELIEVES**

**CHEST CONGESTION**

\*COMPARE TO THE ACTIVE INGREDIENTS OF  
ROBITUSSIN DM®

**ALCOHOL FREE**

**COUGH FORMULA  
FOR ADULTS & CHILDREN**

**16 FL OZ (473 mL)**

### Drug Facts

<b>Active ingredient (per teaspoonful = 5mL)</b>	<b>Purposes</b>
Dextromethorphan HBr, USP 10 mg.....	Cough suppressant
Guaifenesin, USP 100 mg.....	Expectorant

**Uses** • temporarily relieves cough due to minor throat and bronchial irritation • helps loosen phlegm (mucus) • helps thin bronchial secretions to make coughs more productive

**Warnings** • Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Do not use** if you are hypersensitive to any of the ingredients

**Ask a doctor before use if you have**

• A cough with too much phlegm (mucus) • A chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away

**Directions** • take every 4 hours as needed, or as directed by a doctor

• Do not take more than 6 doses in 24 hours

• Do not exceed recommended dose

Adults and children 12 years and over	2 teaspoonfuls (10mL)
Children under 12 years	consult a doctor

**Inactive ingredients:** Citric acid, FD&C Red #40, flavors, glucose, glycerin, high fructose corn syrup, menthol, saccharin sodium, sodium benzoate, water.

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.**

**Other information** • store at room temperature 15°-30°C (59°-86°F)

• protect from freezing • protect from light • for institutional use only

• each teaspoon (5mL) contains: sodium 2 mg

\*This product is not manufactured or distributed by the owner of the registered trademark Robitussin DM®.

Distributed by:

**GERI-CARE  
PHARMACEUTICALS CORP.**  
1650 63rd Street  
Brooklyn, New York, 11204



N  
3 57896-66016 1

Rev. 0404R

## GERI-TUSSIN DM

dextromethorphan hydrobromide and guaifenesin liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-660
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL</b> (UNII: L7T10EP3A)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>WATER</b> (UNII: 059QF0K00R)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-660-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2004	02/28/2021

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/01/2004	02/28/2021

**Labeler** - Geri-Care Pharmaceuticals, Corp (611196254)

**Registrant** - Geri-Care Pharmaceuticals, Corp (611196254)