

**MULTI SYMPTOM SEVERE COLD AND NIGHTTIME SEVERE COLD AND COUGH-
acetaminophen, dextromethorphan hbr, diphenhydramine hcl, phenylephrine
hcl
WALGREEN CO.**

**Walgreens Multi- Symptom Wal-Flu[®] Severe Cold and Nighttime Wal-Flu[®]
Severe Cold and Cough Day & Night Value Pack**

Active ingredients (in each packet) - Multi Symptom Severe Cold

Acetaminophen 500 mg

Dextromethorphan hydrobromide 20 mg

Phenylephrine hydrochloride 10 mg

Active ingredients (in each packet) - Nighttime Severe Cold and Cough

Acetaminophen 650 mg

Diphenhydramine hydrochloride 25 mg

Phenylephrine hydrochloride 10 mg

Purpose - Multi Symptom Severe Cold

Pain reliever/fever reducer

Cough Suppressant

Nasal Decongestant

Purpose - Nighttime Severe Cold and Cough

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

Uses - Multi Symptom Severe Cold

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Uses - Nighttime Severe Cold and Cough

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- runny nose
- sneezing
- itchy nose or throat
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings - Multi Symptom Severe Cold /Nighttime Severe Cold and Cough

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use - Multi Symptom Severe Cold

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not use - Nighttime Severe Cold and Cough

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have - Multi Symptom Severe Cold

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor before use if you have - Nighttime Severe Cold and Cough

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are - Multi Symptom Severe Cold

taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are - Nighttime Severe Cold and Cough

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product - Multi Symptom Severe Cold

do not exceed recommended dosage

When using this product - Nighttime Severe Cold and Cough

- **do not exceed recommended dosage**
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if - Multi Symptom Severe Cold /Nighttime Severe Cold and Cough

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions - Multi Symptom Severe Cold

- **do not use more than directed**
- take every 4 hours, while symptoms persist .do not take more than 6 packets in 24 hours unless directed by a doctor

Age	Dose
Adults and children 12 years of age	One packet

Children under 12 years of age

Do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10 - 15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating, Do not overheat

Directions Nighttime Severe Cold and Cough

- **do not use more than directed**
- take every 4 hours while symptoms persist, not to exceed 5 packets in 24 hours unless directed by a doctor

Age	Dose
Adults and Children 12 years of age and over	One packet
Children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10 - 15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating, Do not overheat.

Other information - Multi Symptom Severe Cold

- each packet contains: **potassium 6 mg**
- store at room temperature. Protect from excessive heat and moisture.

Other information - Nighttime Severe Cold and Cough

- each packet contains: **potassium 6 mg**
- store at room temperature. Protect from excessive heat and moisture

Inactive ingredients - Multi Symptom Severe Cold

citric acid, caramel, flavors, maltodextrin, potassium chloride, silica ,sucralose, sucrose

Inactive ingredients - Nighttime Severe Cold and Cough

citric acid, F D & C yellow #6 , flavors, maltodextrin, potassium chloride, silica ,sucralose, sucrose,

Questions or comments?

1-866-467-2748

Package/Label Principal Display Panel

Walgreens

DAY & NIGHT PACK

Compare to Theraflu® Multi Symptom Severe Cold and Nighttime Severe Cold & Cough active ingredients††

MULTI- SYMPTOM

Wal-Flu®

Severe

COLD

ACETAMINOPHEN / PAIN RELIEVER/ FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

ASPARTAME FREE

SODIUM FREE

- Relieves nasal congestion, sore throat pain, cough, headache, body ache & fever

GREEN TEA & HONEY LEMON FLAVORS

INFUSED WITH MENTHOL AND GREEN TEA FLAVOR

6 PACKETS

NIGHTTIME

Wal-Flu®

Severe

COLD & COUGH

ACETAMINOPHEN/ PAIN RELIEVER/ FEVER REDUCER

DIPHENHYDRAMINE HCl / ANTIHISTAMINE/ COUGH SUPPRESSANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

ASPARTAME FREE

SODIUM FREE

Relieves nasal congestion, sore throat pain, cough, headache, body ache & fever, runny nose

& sneezing

HONEY LEMON INFUSED WITH CHAMOMILE & WHITE TEA FLAVORS

6 PACKETS

TOTAL 12 PACKETS

**READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE .KEEP
CARTON FOR REFERENCE, DO NOT DISCARD.**

**DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5
DOSES IN TOTAL IN ANY 24 -HOUR PERIOD.**

**DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS
AFTER THE LAST DOSE OF DAYTIME PRODUCT UNLESS DIRECTED BY
DOCTOR.**

††These Products are not manufactured or distributed by GSK Consumer Healthcare,
distributor of Theraflu[®] Daytime Severe Cold & Cough and Theraflu[®] Nighttime Severe
Cold & Cough

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Walgreens 100% SATISFACTION GUARANTEED

Walgreens.com ©2018 Walgreen Co,

WALGREENS PHARMACIST RECOMMENDED‡

Walgreens Pharmacist Survey

**TAMPER EVIDENT INNER UNIT:
DO NOT USE IF SEALED PACKET IS TORN OR BROKEN**



ORG code is an internal Periscope code, that will be updated by Periscope prior to network release. It is subject to change.

MULTI-SYMPOM SEVERE COLD AND NIGHTTIME SEVERE COLD AND COUGH

acetaminophen, dextromethorphan hbr, diphenhydramine hcl, phenylephrine hcl kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-5480

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5480-12	1 in 1 CARTON; Type 0: Not a Combination Product	06/22/2018	

Quantity of Parts			
Part #	Package Quantity	Total Product Quantity	
Part 1	6 PACKET	6	
Part 2	6 PACKET	6	

Part 1 of 2

MULTI SYMPTOM SEVERE COLD MULTI SYMPTOM SEVERE COLD

acetaminophen, dextromethorphan hbr, phenylephrine hcl powder, for solution

Product Information

Item Code (Source) NDC:0363-5479

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARAMEL (UNII: T9D99G2B1R)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5479-06	6 in 1 CARTON		
1		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/31/2018	

Part 2 of 2

NIGHTTIME SEVERE COLD AND COUGH NIGHTTIME SEVERE COLD AND COUGH

acetaminophen, diphenhydramine hcl, phenylephrine hcl powder, for solution

Product Information

Item Code (Source) NDC:0363-5481

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SUCROSE (UNII: C151H8M554)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5481-06	6 in 1 CARTON		
1		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/29/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M013	06/22/2018	
--------------------	------	------------	--

Labeler - WALGREEN CO. (008965063)

Revised: 11/2023

WALGREEN CO.