DAYTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated DISCOUNT DRUG MART

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1178-DDM-2022-0809

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion and pressure
 - cough due to minor throat & bronchial irritation
 - minor aches and pains
 - headache
 - fever
 - sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by

fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

take only as directed (see overdose warning)

do not exceed 4 doses per 24 hours

adults and children 12 years and over	take 2 caplets every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

DISCOUNT drug mart FOOD MARKET

NDC 53943-178-03

†Compare to the active ingredients in Vicks $\ \ \$ DayQuil $\ \ \ \ \$ Severe Cold $\ \ \$ Flu Caplet

DAYTIME SEVERE COLD & FLU

MAXIMUM STRENGTH

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guaifenesin

Pain Reliever/Fever Reudcer, Nasal Decongestant, Cough Suppressant, Expectorant

Headache, Fever, Sore Throat, Minor Aches & Pains


Nasal/Sinus Congestion & Sinus Pressure

Cough

Chest Congestion

Actual Size

24 CAPLETS



Distributed by: Drug Mart - Food Fair 211 Commerce Drive, Medina, OH 44256

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

estions or comments?

 thylene glycol, polyvinyl alcohol, povidone, pregelatinized
 h, stearic acid, talc, titanium dioxide nesium stearate, maltodextrin, microcrystalline cellulose *ug Facts* **(**continued)

Drug Facts

ann fine
Active ingredients (in each caplet)
Acetaminophen 325 mg Pain reliever/fever red
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephine HCl 5 mgNasal deconge

sinus congestion and press ■ Sore headache minor aches and pains temporarily relieves common cold/flu symptoms: cough due to minor throat & bronchial irritation

restores freer breathing through the nose

promotes rasal and/or sinus dra helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus reduces swelling of nasal passages

make coughs more productive

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters 3 or more alcoholic drinks every day while using this product more than 8 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning**: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have Inter disease Inter disease International blood pressure International disease Indian for the formuch philegm (mucus) Internating due to an enlarged prostate gland International Cough that occurs with too much philegm (mucus)

persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin When using this product do not use more than directed.

Stop use and ask a doctor if

fever gets worse or lasts more than 3 Stop use and ask a doctor if wou get nervous, dizzy, or sleepless pain, nasal congestion, or cough gets worse or lasts more than 7 days

 cough comes back or occurs with rash or headache that new symptoms occur These could be signs of a serious condition. redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical h contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms

This product is not manufactured or distributed by Proctor & Gamble, distributor of Vicks® DayQuil® Severe Cold & Flu.

F117803DDM_R1

4

015

9332,

0



TIME SEVER

MAXIMUM STRENGTH



NDC 53943-178-03

†Compare to the active ingredients in Vicks* Dayquil* Severe Cold & Flu

ME

MAXIMUM STRENGTH

Acetaminophen, Phenylephrine HCI, Dextromethorphan HBr, Guaifenesin Pain Reliever / Fever Reducer, Nasal Decongestant, Cough Suppressant, Expectorant

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal/Sinus Congestion & Sinus Pressure

Cough

Chest Congestion



24 CAPLETS

DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product	Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:53943-178

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)			
CROSPOVIDONE (UNII: 2S7830E561)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POLYVINYL ALCOHOL (UNII: 532B59J990)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
STARCH, CORN (UNII: O8232NY3SJ)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	AAA;1178
Contains			
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:53943- 178-03	2 in 1 CARTON	08/18/2014			
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/18/2014	

Labeler - DISCOUNT DRUG MART (047741335)

Revised: 8/2022 DISCOUNT DRUG MART