

**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled**  
**CVS Pharmacy**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®†  
Cold/Flu

### **Multi Symptom**

Nighttime

### **Cold/Flu Relief**

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

DOXYLAMINE SUCCINATE - Antihistamine

Relieves:

Aching, fever, sore throat, coughing, sniffing and sneezing so you can rest

SOFTGELS\*\*

(\*\*LIQUID-FILLED CAPSULES)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademark of the Procter & Gamble

Company

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by: CVS Pharmacy, Inc

One CVS Drive, Woonsocket, RI 02895

CVS.com

**Package Label**

**CVS Health**

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®†

**Cold/Flu**

NDC 69842-847-16

**Multi-Symptom Nighttime Cold/Flu Relief**

**ACETAMINOPHEN** - Pain reliever; Fever reducer  
**DEXTROMETHORPHAN HBr** - Cough suppressant  
**DOXYLAMINE SUCCINATE** - Antihistamine

**Relieves:**  
 Aching, fever, sore throat, coughing, sniffing, and sneezing so you can rest

**16 SOFTGELS\*\*** (LIQUID-FILLED CAPSULES)

**Drug Facts**

**Active ingredients (in each softgel)**

- Acetaminophen 325 mg.....Pain reliever/fever reducer
- Dextromethorphan HBr 15 mg.....Cough suppressant
- Doxylamine succinate 6.25 mg.....Antihistamine

**Purposes**

- temporarily relieves common cold and flu symptoms: sore throat, headache, minor aches and pains, fever, runny nose and sneezing
- cough due to minor throat and bronchial irritation

**Uses**

- temporarily relieves common cold and flu symptoms: sore throat, headache, minor aches and pains, fever, runny nose and sneezing
- cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours; with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening; blisters; rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** liver disease; glaucoma; trouble urinating due to an enlarged prostate gland; a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema; cough that occurs with too much phlegm (mucus).

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin; taking sedatives or tranquilizers.

**Drug Facts (continued)**

**When using this product** do not exceed recommended dosage; excitability may occur, especially in children; avoid alcoholic drinks; marked drowsiness may occur; alcohol, sedatives, and tranquilizers may increase drowsiness; be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if** pain or cough gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** do not take more than directed (see Overdose warning); do not take more than 4 doses in 24 hours; adults and children 12 years and over: take 2 softgels with water every 6 hours; swallow whole; do not crush, chew, or dissolve; children under 12 years: do not use; when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing.

**Other information** store between 15-30°C (59-86°F); avoid excessive heat.

**Inactive ingredients** D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink.

**Questions or comments?**  
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Lot No.:  
 Exp. Date:

**CVS HEALTH Nighttime Cold and Flu Relief**

**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-847
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69842-847-16	16 in 1 CARTON	03/31/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69842-847-24	24 in 1 CARTON	03/31/2019	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:69842-847-48	48 in 1 CARTON	03/31/2019	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
---------------------------	---	-----------------------------	---------------------------

OTC Monograph Drug M012

03/31/2019

**Labeler** - CVS Pharmacy (062312574)

Revised: 10/2024

CVS Pharmacy