

TYLENOL PM EXTRA STRENGTH- acetaminophen and diphenhydramine hydrochloride tablet, film coated
Morning Star OTC

Tylenol® PM Extra Strength

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 500 mg	Pain reliever
Diphenhydramine HCl 25 mg	Nighttime sleep aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets at bedtime▪ do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use

Other information

- store between 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing**

Inactive ingredients

carnauba wax, crospovidone, FD&C blue no. 1 aluminum lake, hypromellose, magnesium

Drug Facts (continued)

Inactive ingredients: camphor wax, croscopollose, FD&C blue no. 1, aluminum lake, hydroxyethyl magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 85, polyvinyl, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

TYLENOL Extra Strength **2 CAPLETS**

TYLENOL Extra Strength **2 CAPLETS**

Acetaminophen **diphenhydramine HCl**

Pain Reliever, Nighttime Sleep Aid Non-habit forming

2 CAPLETS

Morning Star OTC

DO NOT USE IF POUCH IS TORN OR DAMAGED

Drug Facts

Active ingredients (in each caplet)

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain Reliever
Diphenhydramine HCl 25 mg	Nighttime Sleep Aid

Uses
Temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness.

Warnings
Lower Warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen.
 ■ 8 or more alcoholic drinks a day while taking this product.
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin rash including:
 ■ hives ■ rash
 ■ skin redness
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other product containing diphenhydramine, even one used on skin.
 ■ in children under 12 years of age.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have
 ■ liver disease
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland
 ■ diabetes

LEFT PANEL FOR CONTINUED DRUG FACTS

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 ©J&JCI 2020

This product is repackaged by:
Morning Star OTC
 145 S. Anderson St., Los Angeles, CA 90033
 Hours of operation 9 am - 4 pm, 323.354.4838



LOT
EXP

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin.
 ■ taking sedatives or tranquilizers.

When using this product
 ■ drowsiness will occur
 ■ do not drink a motor vehicle or operate machinery.

Stop use and ask a doctor if
 ■ sleeplessness persists continuously for more than 2 weeks
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur.

These could be signs of a serious condition.
 ■ pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ do not take more than directed (see overdose warning)
adults and children 12 years and older
 • take 2 caplets at bedtime
children under 12 years
 • do not take more than 2 caplets of this product in 24 hours
 • do not use.

Other information
 ■ Use between 20-32°C (68-77°F)
 ■ do not use if pouch is torn or damaged



TYLENOL PM EXTRA STRENGTH

acetaminophen and diphenhydramine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53209-1002(NDC:50580-608)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue (Light Blue)	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TY;PM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53209-1002-2	25 in 1 POUCH	06/19/2025	
1	NDC:53209-1002-1	2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/19/2025	

Labeler - Morning Star OTC (078589357)

Registrant - Morning Star OTC (078589357)

Establishment

Name	Address	ID/FEI	Business Operations
Morning Star OTC		078589357	repack(53209-1002)