

BACK AND BODY EXTRA STRENGTH- aspirin, caffeine tablet, film coated
L.N.K. International, Inc.

Quality Plus 44-413

Active ingredients (in each caplet)

Aspirin 500 mg (NSAID)*

Caffeine 32.5 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever aid

Uses

for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have asthma
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

Questions or comments?

1-800-426-9391

Principal display panel

NDC 50844-413-08

†Compare to active ingredients in
Bayer® Back & Body

**QUALITY
+PLUS**

EXTRA STRENGTH

BACK & BODY

Aspirin (NSAID) and Caffeine
PAIN RELIEVER/PAIN RELIEVER AID

24

Coated Caplets

Relieves Backaches &
Muscle Aches & Pains

ACTUAL
SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed
by Bayer AG, owner of the registered trademark
Bayer® Back & Body.
50844 REV0222A41308

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

B-1603-413-08
REV0222A41308

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Aspirin 500 mg (NSAID)* Pain reliever
Caffeine 32.5 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of

- backache pain
- muscle aches and pains
- minor aches and pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product

*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Back & Body.
50844 REV0222A41308
Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

QUALITY PLUS

NDC 50844-413-08

Compare to active ingredients in Bayer® Back & Body

EXTRA STRENGTH

BACK & BODY

Aspirin (NSAID) and Caffeine
PAIN RELIEVER/PAIN RELIEVER AID

24 Coated Caplets

Relieves Backaches & Muscle Aches & Pains

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



No Print Area
Lot no. & Exp. Date

Drug Facts (continued)

- new symptoms occur
- These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions do not take more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, hypromellose, methoxyfen, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate

Questions or comments?
1-800-426-9391

Quality Plus 44-413

BACK AND BODY EXTRA STRENGTH

aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-413
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	32.5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	44;413
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-413-09	1 in 1 CARTON	04/12/2004	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:50844-413-08	1 in 1 CARTON	04/12/2004	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/12/2004	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-413)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(50844-413)
-------------------------	--	-----------	------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-413)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-413)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-413)

Revised: 9/2025

L.N.K. International, Inc.