

**BMD 10%- bacitracin methylene disalicylate granule**  
**Phibro Animal Health**

-----  
**BMD 10%**

BMD 10%

| <b>Species for Use</b> | <b>Dose</b> | <b>Indications</b> |
|------------------------|-------------|--------------------|
|------------------------|-------------|--------------------|

|   |        |                       |
|---|--------|-----------------------|
| <b>Broiler and replacement chickens:</b><br>weight gain and improved feed efficiency. | 4 - 50 | For increased rate of |
|---|--------|-----------------------|

|   |    |                            |
|---|----|----------------------------|
| necrotic enteritis caused or complicated by Clostridium spp. or other organisms susceptible to bacitracin methylene disalicylate. | 50 | As an aid in prevention of |
|---|----|----------------------------|

|   |         |   |
|---|---------|---|
| necrotic enteritis caused or complicated by organisms susceptible to bacitracin | 100-200 | As an aid in control of Clostridium spp. or other methylene disalicylate. |
|---|---------|---|

|   |  |   |
|---|--|---|
| <b>CAUTION:</b> To control a necrotic enteritis outbreak, start medication disease. The dosage range permitted levels based on severity of the infection. diagnostic laboratory or pathologist to determine regarding the optimal level of drug. for 5 - 7 days or as long as clinical signs medication to prevention level (50 g/ton). |  | at the first clinical signs of provides for different Consult a poultry the diagnosis and advice Administer continuously persist, and then reduce |
|---|--|---|

|  |         |   |
|--|---------|---|
| <b>Laying Hens:</b><br>production and improved feed efficiency during production | 10 - 25 | For increased egg the first seven months of |
|--|---------|---|

|  |        |  |
|--|--------|--|
| <b>Growing turkeys:</b><br>of weight gain and improved feed efficiency.        | 4 - 50 | For increased rate   |
| transmissible enteritis in growing turkeys susceptible to bacitracin methylene | 200    | As an aid in control of complicated by organisms disalicylate. |

|                   |        |                    |
|-------------------|--------|--------------------|
| <b>Pheasants:</b> | 4 - 50 | For increased rate |
|-------------------|--------|--------------------|

of weight gain and improved feed efficiency.

**Quail:**

of weight gain and improved feed efficiency in age.

5 - 20

For increased rate

quail not over 5 weeks of

200  
enteritis in growing quail due to  
susceptible to bacitracin methylene disalicylate.  
sole ration.

For prevention of ulcerative  
*Clostridium colinum*  
Feed continuously as the

**Growing/Finishing Swine:**

weight gain and improved feed efficiency.

10 - 30

For increased rate of

250  
dysentery (bloody scours) associated with  
pigs up to 250 lbs body weight. Feed  
complete feed on premises with a history of  
signs of the disease have not yet  
approved treatment of the disease

For control of swine  
*Brachyspira hyodysenteriae* in  
250 grams per ton of  
swine dysentery, but where  
occurred or following an  
condition.

5 to 7 mg/lb in swine weighing 40 to  
confirmed by a veterinarian when

The 250 g/ton level will provide  
250 lbs.

Feed containing an approved level of  
disalicylate should be the sole ration.

CAUTION: Diagnosis should be  
results are not satisfactory.  
bacitracin methylene

**Pregnant Sows:**

clostridial enteritis caused by *C. perfringens* in  
sows from 14 days before through 21  
premises with a history of clostridial scours.

250

For control of

mg/head/day  
confirmed by a veterinarian when

suckling piglets when fed to  
days after farrowing on

Feed containing an approved level of  
disalicylate should be the sole ration.

CAUTION: Diagnosis should be  
results are not satisfactory.  
bacitracin methylene

**Feedlot beef cattle:**  
condemnations due to  
cattle.

70 For reduction in the number of liver  
(continuously) abscesses in feedlot beef

liver condemnations due to  
cattle.

250 For reduction in the number of  
(continuously) abscesses in feedlot beef

Note: Where minimum levels are shown, increase the antibiotic concentration within approved range when necessary to fit the feeding program, and to insure adequate levels of antibiotic in the complete ration.

BMD 10 Label



## BMD 10%

bacitracin methylene disalicylate granule

### Product Information

**Product Type**

OTC ANIMAL DRUG

**Item Code (Source)**

NDC:66104-6409

**Route of Administration** ORAL

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength           |
|---|-------------------|--------------------|
| <b>BACITRACIN METHYLENEDISALICYLATE</b> (UNII: JGV6OJ52FT) (BACITRACIN - UNII:58H6RWO52I) | BACITRACIN        | 50 g<br>in 0.45 kg |

### Product Characteristics

|                 |                      |                     |  |
|-----------------|----------------------|---------------------|--|
| <b>Color</b>    | gray (Gray to Brown) | <b>Score</b>        |  |
| <b>Shape</b>    |                      | <b>Size</b>         |  |
| <b>Flavor</b>   |                      | <b>Imprint Code</b> |  |
| <b>Contains</b> |                      |                     |  |

### Packaging

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:66104-6409-1 | 50 kg in 1 BAG      |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| export only        |  | 02/19/2025           |                    |

**Labeler** - Phibro Animal Health (006989008)

**Registrant** - Phibro Animal Health (006989008)

Revised: 2/2025

Phibro Animal Health