

**NAPROXEN SODIUM- naproxen sodium tablet**  
**Bi-Mart**

-----

**ACTIVE INGREDIENT**

**(in each caplet)**

Naproxen sodium 220 mg (Naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**PURPOSE**

Pain reliever/fever reducer

**USES**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**WARNINGS**

**Allergy alert:**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**DO NOT USE**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**ASK A DOCTOR BEFORE USE IF**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefits of aspirin
- taking any other drug

**WHEN USING THIS PRODUCT**

- take with food or milk if stomach upset occurs

**STOP USE AND ASK DOCTOR IF**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

## **PREGNANCY/BREASTFEEDING**

**If pregnant or breast-feeding**,ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## **DIRECTIONS**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

<b>adults and children 12 years and older</b>	<ul style="list-style-type: none"> <li>• take 1 caplet every 8 to 12 hours while symptoms last</li> <li>• for the first dose you may take 2 caplets within the first hour</li> <li>• do not exceed 2 caplets in any 8 to 12 hour period</li> <li>• do not exceed 3 caplets in a 24 hour period</li> </ul>
<b>children under 12 years</b>	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

## **OTHER INFORMATION**

- **each caplet contains:**sodium 20 mg
- store at 20 - 25°C (68 - 77°F).
- avoid high humidity and excessive heat above 40°C (104°F)
- DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

## **INACTIVE INGREDIENTS**

FD&C blue#2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide

## **QUESTIONS?**

call toll free **1-844-912-4012**

Eugene, OR 97402

**NDC 37835-594-01**

## 100 Caplets

\*\*This product is not manufactured or distributed by the owner of the registered trademark ALEVE®.

0 713571 903481



1

**BI-MART** NDC 37835-594-01

Compare to the active ingredient in ALEVE<sup>®</sup>

**FAST LONG-LASTING PAIN RELIEF**

**NAPROXEN SODIUM**

CAPLETS 220 MG



**100 CAPLETS**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Distributed by:  
**BI-MART**  
Eugene, OR 97402

**Drug Facts**

**Active ingredient (in each caplet)** *Purpose*  
Naproxen sodium 220 mg (NSAID)\* *Pain reliever/fever reducer*  
(naproxen 220 mg) (mgi) (NSAID)\* *Nonsteroidal anti-inflammatory drug*

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis ■ muscular aches ■ backache
- menstrual cramps ■ headache ■ toothache ■ the common cold ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

PEEL FOR DIRECTIONS

**171 FINE**

594-01-BM 02/25

**Drug Facts (continued)**

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if** you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center: (1-800-222-1222) right away.

**Directions**

- do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose
- adults and children 12 years and older: take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8 to 12-hour period ■ do not exceed 3 caplets in a 24-hour period
- children under 12 years: ask a doctor

**Other information** ■ each caplet contains: sodium 20 mg ■ store at 20° - 25°C (68° - 77° F) ■ avoid high humidity and excessive heat above 40°C (104°F)

**Inactive ingredients** FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, powdered k-30, sodium starch glycolate, stearic acid, titanium dioxide.

**Questions?** call toll-free 1-844-912-4012

\*\*This product is not manufactured or distributed by the owner of the registered trademark *Advil*®

**MARKET INFORMATION: ADVIL ASK GUARANTEES**  
If you are not completely satisfied with your **Advil**® purchase, please return it for a full refund. No receipt necessary. **Advil** is a registered trademark of **Pfizer Inc.** © 2005 **Pfizer Inc.**

**NDC 37835-594-03**

NAPROXEN SODIUM

CAPLETS 220 MG  
PAIN RELIEVER/FEVER REDUCER (NSAID)  
WITH 8 TO 12 HOUR DOSING  
300 Caplets

**\*\*This product is not manufactured or distributed by the owner of the registered trademark ALEVE®.**

VP  
VALUE  
PACK

**BI-MART**  
FAST LONG-LASTING PAIN RELIEF  
**NAPROXEN  
SODIUM**  
CAPLETS 220 MG

**EASY OPEN**  
SENIOR FRIENDLY CAP  
**300 CAPLETS**

0 73579 03031 8

Compare to the active ingredient in  
**ALEVE®**  
(naproxen 200 mg) (NSAID)\*

**PAIN RELIEVER/FEVER REDUCER (NSAID)  
WITH 8 TO 12 HOUR DOSING**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active ingredient (in each caplet)** *Purposes*  
Naproxen sodium 220 mg.....Pain reliever/fever reducer  
(naproxen 200 mg) (NSAID)\*  
*\*nonsteroidal anti-inflammatory drug*

**Uses**  
temporarily relieves minor aches and pains due to:  
■ minor pain of arthritis ■ muscular aches ■ backache  
■ menstrual cramps ■ headache ■ toothache ■ the common  
cold ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic  
reaction, especially in people allergic to aspirin. Symptoms may  
include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right  
away.  
**Stomach bleeding warning:** This product contains an NSAID,  
which may cause severe stomach bleeding. The chance is higher  
if you: ■ are age 60 or older ■ have had stomach ulcers or  
bleeding problems ■ take a blood thinning (anticoagulant) or  
steroid drug ■ take other drugs containing prescription or  
nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this  
product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin,  
increase the risk of heart attack, heart failure, and stroke. These  
risks are higher if you use more than directed or for  
longer than directed.  
**Do not use** ■ if you have ever had an allergic reaction to any other  
pain reliever/fever reducer ■ right before or after heart surgery  
**Ask a doctor before use if** ■ the stomach bleeding warning

**PEEL FOR DIRECTIONS**  
594-03-BM 03/25  
**LIFT HERE**

**Drug Facts (continued)**  
applies to you ■ you have a history of stomach problems, such  
as heartburn ■ you have high blood pressure, heart disease,  
liver cirrhosis, kidney disease, asthma, or had a stroke ■ you  
are taking a diuretic ■ you have problems or serious side  
effects from taking pain relievers or fever reducers  
**Ask a doctor or pharmacist before use if you are** ■ under a  
doctor's care for any serious condition ■ taking aspirin for heart  
attack or stroke, because naproxen may decrease this benefit of  
aspirin ■ taking any other drug  
**When using this product** ■ take with food or milk if stomach  
upset occurs  
**Stop use and ask a doctor if** ■ you experience any of the  
following signs of stomach bleeding: ■ red faint ■ vomit blood  
■ have bloody or black stools ■ have stomach pain that does  
not get better ■ you have symptoms of heart problems or  
stroke: ■ chest pain ■ trouble breathing ■ weakness in one part  
or side of body ■ slurred speech ■ leg swelling ■ pain gets  
worse or lasts more than 10 days ■ fever gets worse or lasts  
more than 3 days ■ you have difficulty swallowing ■ it feels  
like the pill is stuck in your throat ■ redness or swelling is  
present in the painful area ■ any new symptoms appear  
**If pregnant or breast-feeding,** ask a health professional before  
use. It is especially important not to use naproxen sodium at 20  
weeks or later in pregnancy unless definitely directed to do so  
by a doctor because it may cause problems in the unborn child  
or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical  
help or contact a Poison Control Center (1-800-222-1222) right  
away.

**Directions**  
■ do not take more than directed  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose  
**Adults and  
children 12  
years and older**  
■ take 1 caplet every 8 to 12 hours while  
symptoms last ■ for the first dose you may  
take 2 caplets within the first hour ■ do not  
exceed 2 caplets in any 8 to 12-hour period  
■ do not exceed 3 caplets in a 24-hour period  
■ ask a doctor  
**children under  
12 years**  
**Other information** ■ each caplet contains: sodium 20 mg  
■ store at 20°-25° C (68°-77° F) ■ avoid high humidity and excessive  
heat above 40° C (104° F)  
**Inactive ingredients** FD&C blue #2 aluminum lake,  
hypromellose 2810, maize starch, microcrystalline cellulose,  
polyethylene glycol, polydione K-30, sodium starch glycolate,  
stearic acid, titanium dioxide.  
**Questions?** call toll-free 1-844-912-4012  
\*\*This product is not manufactured or distributed by the owner of the  
registered trademark Aleve®  
Distributed by:  
**BI-MART**, Eugene, OR 97402  
EVEN IF THE PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY THE OWNER OF THE  
REGISTERED TRADEMARK ALEVE®, IT MAY BE COVERED BY A PATENT.

NAPROXEN SODIUM			
naproxen sodium tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-594
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg
Inactive Ingredients			
Ingredient Name			Strength
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			

STARCH, CORN (UNII: O8232NY3SJ)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE K30 (UNII: U725QWY32X)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	blue (Light Blue)		Score	no score
Shape	OVAL (Caplet -Shaped)		Size	12mm
Flavor			Imprint Code	220
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-594-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:37835-594-03	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA091353		09/30/2011	

**Labeler -** Bi-Mart (027630078)

**Establishment**

Name	Address	ID/FEI	Business Operations
Granules India Ltd		918609236	manufacture(37835-594)