# CARE ONE ASPIRIN- as pirin tablet, chewable American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### American Sales Company Aspirin Drug Facts

### **Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

### Purpose

Pain reliever

### Uses

for the temporary relief of minor aches and pains

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

#### Other information

• store at 20-25°C (68-77°F)

### **Inactive ingredients**

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

### Questions or comments?

1-800-719-9260

### **Principal Display Panel**

Compare to Active Ingredient in Bayer® Chewable Aspirin

Adult Low Strength

**ASPIRIN** 

81 mg

**Actual Size** 

Pain Reliever (NSAID)

Chewable Orange Flavor

OUR PHARMACISTS RECOMMEND

Low Dose Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

36 TABLETS



46768 OF C2

### CODE VIBER



CAP IS BROKEN OR MISSING DO NOT USE IF PRINTED SEAL UNDER

#### Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)\*..... ..Pain reliever \*nonsteroidal anti-inflammatory drug

Purpose

Uses for the temporary relief of minor aches and pains

#### Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling

- ∎asthrna (wheezing) shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higherifyou ■ are age 60 or older
- have had stomach ulcers or bleeding. problems
- take a blood thinning (anticoagulant) or steroid drug

#### Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use ■if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ for at least 7 days after ton sill ectomy or oral surgery unless directed by a doctor

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney diseas
- you are taking a diuretic you have asthma Ask a doctor or p harma cist before use if you are taking a prescription drug for

#### ■ gout ■ diabetes ■ arthritis

#### Stop use and ask a doctor if

- an aller gic reaction occurs. Seek medical help
- right away. you experience any of the foll owing signs of stomach bleeding:
- feel faint womit blood
- have bloody or black stools
- have sto mach pain that does not get better ▶

#### Drug Facts (continued)

- pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symp toms occur
- ringing in the ears or aloss of hearing occurs These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

#### Directions

- drink a full glass of water with each close
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

### Other information

■ store at 20-25°C(68-77°F)

Gluten Free
This poductisnot manufactued or distributed by Baye
Hastinicase LLC, distributor of Bayero Charactue, sprin, Qually DISTRIBUTED BY FOODHOLD U.S.A., LLC LANDOVER, MD 20785-1-877-946-9949 92018 S86 Brands, LLC Inactive ingredients com starch, dex trose excipient, FD&C yellow no. 6 aluminum lake, flavors, sæcharin sodium Drug Questions or comments? 1-800-719-9260 guaranteed Facts (continued) or your money back

### **CARE ONE ASPIRIN**

aspirin tablet, chewable

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-467
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	

Inactive Ingredients		
Ingredient Name	Strength	
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)		
STARCH, CORN (UNII: O8232NY3SJ)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		

Product Characteristics			
Color	ORANGE	Score	no score
Shape	ROUND	Size	8 mm
Flavor	ORANGE	Imprint Code	L467
Contains			

1	Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date	
1	NDC:41520-467-68	1 in 1 CARTON	06/01/2004		
1		36 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/01/2004	

## Labeler - American Sales Company (809183973)

Revised: 8/2018 American Sales Company