

NICOTINE TRANSDERMAL SYSTEM- nicotine transdermal system patch
Aveva Drug Delivery Systems, Inc.

Nicotine Transdermal System

Drug Facts

Active ingredient (in each patch)

Nicotine, 21 mg delivered over 24 hours

Active ingredient (in each patch)

Nicotine, 14 mg delivered over 24 hours

Active ingredient (in each patch)

Nicotine, 7 mg delivered over 24 hours

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances, remove the patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- **If you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed self-help guide for complete directions and other information.
- begin using the patch on your quit day
- if you smoke **more than 10 cigarettes per day**, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
Step 1 use one 21 mg patch/day	Step 2 use one 14 mg patch/day	Step 3 use one 7 mg patch/day

- if you smoke **10 or less cigarettes per day**, start with **Step 2** for 6 weeks, then **Step 3** for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean, and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces

- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information

- store at 20°C - 25°C (68°F - 77°F)

Inactive ingredients

acrylate adhesive, polyester, silicone adhesive

Questions or comments?

1-888-889-8893

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

For your family's protection, patches are supplied in child-resistant pouches.

Do not use if individual pouch is open or torn.

See self-help guide inside.

*Quitine Circle™ is an optional community support program and is not related to the therapeutic effect, treatment duration, or clinical outcomes of this product.

Distributed by:
QUITINE LLC
Lewes, DE, 19958

Rev. 12/2025

PRINCIPAL DISPLAY PANEL - STEP ONE 21 mg 14 CARTON

NDC 3215-1206-14

Quitine

NICOTINE TRANSDERMAL SYSTEM PATCH

STOP SMOKING AID

STEP ONE

21 mg

14 CLEAR PATCHES

2 week kit

If you smoke **MORE** than 10 cigarettes per day: Start with Step 1.

If you smoke 10 or **LESS** cigarettes per day: Start with Step 2.

Free access to the Quitine Circle™*

A supportive quite community

21 mg delivered over 24 hours

QUITTING IS HARD. YOU DON'T HAVE TO DO IT ALONE.

Join the Quitine Circle™
A free support community for quitting smoking or vaping.

Scan to get instant support

NDC 3215-1206-14

Quitine
NICOTINE TRANSDERMAL SYSTEM
PATCH
STOP SMOKING AID

STEP ONE
21 mg
14 CLEAR PATCHES
2 week kit

SCAN TO SPIN. EVERY SCAN WINS.
*Takes 10 seconds. Win gift cards, big discounts, or even a year's supply of Quitine.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale in vending machines or from any source where proof of age cannot be verified.

See call-help guide inside.

*Quitine Circle™ is an optional community support program and is not related to the therapeutic effect, treatment duration, or clinical outcomes of the product.

Distributed by: QUITINE LLC
Lenex, NC, 28859

1004490 Rev. 10/2025

60010-55994

Drug Facts

Active Ingredient (in each patch)
Nicotine, 21 mg delivered over 24 hours

Purpose
Stop smoking aid

Use
reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have:

- Heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- High blood pressure not controlled with medicine. Nicotine can increase your blood pressure.
- An allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- Stomach ulcer or diabetes
- History of seizures

Ask a doctor or pharmacist before use if you are:

- Using a non-nicotine stop smoking drug
- Taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product if you have vivid dreams or other sleep disturbances, remove the patch at bedtime.

Stop use and ask a doctor if:

- Skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- Irregular heartbeat or palpitations occur
- You get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness, and rapid heartbeat
- You have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

DIRECTIONS If you are under 18 years of age, ask a doctor before use. Before using this product, read the enclosed call-help guide for complete directions and other information. Begin using the patch on your quit day. If you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
Step 1 use one 21 mg patch/day	Step 2 use one 14 mg patch/day	Step 3 use one 7 mg patch/day

If you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks. Apply one new patch every 24 hours on skin that is dry, clean, and hairless. Remove backing from patch and immediately press onto skin. Hold for 10 seconds. Smooth hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch. If the used patch should be removed and a new one applied to a different skin site at the same time each day. If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning. Do not wear more than one patch at a time. Do not cut patch in half or into smaller pieces. Do not reuse patch as for more than 24 hours because it may irritate your skin and lose strength after 24 hours. It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other Information Store at 20°C - 25°C (68°F - 77°F)

Inactive Ingredients acrylic adhesive, polyester, silicone adhesive

Questions or comments? 1-800-888-9888

PRINCIPAL DISPLAY PANEL - STEP TWO 14 mg CARTON

NDC 3215-1205-14

Quitine
NICOTINE TRANSDERMAL SYSTEM PATCH
STOP SMOKING AID
STEP TWO

14 mg

14 CLEAR PATCHES

2 week kit

If you smoke MORE than 10 cigarettes per day: Use after completing Step 1.

If you smoke 10 or LESS cigarettes per day: Start with Step 2.

Free access to the Quitine Circle™*

A supportive quite community

14 mg delivered over 24 hours



PRINCIPAL DISPLAY PANEL - STEP THREE 7 mg CARTON

NDC 3215-1204-14

Quitine

NICOTINE TRANSDERMAL SYSTEM PATCH

STOP SMOKING AID

STEP THREE

7 mg

14 CLEAR PATCHES

2 week kit

For use after completing Step 2.

Free access to the Quitline Circle™*

A supportive quitte community

7 mg delivered over 24 hours



NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:3215-1206
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	21 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:3215-1206-14	14 in 1 CARTON	02/18/2026	
1		1 h in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:3215-1206-28	28 in 1 CARTON	02/18/2026	
2		1 h in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074612	02/18/2026	

NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:3215-1205
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	14 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:3215-1205-14	14 in 1 CARTON	02/18/2026	
1		1 h in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074612	02/18/2026	

NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:3215-1204
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	7 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:3215-1204-14	14 in 1 CARTON	02/18/2026	
1		1 h in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074612	02/18/2026	

Revised: 2/2026

Aveva Drug Delivery Systems, Inc.