

BISMUTH SUBSALICYLATE- bismuth subsalicylate tablet, film coated
CVS Pharmacy, Inc.

5346-CVS

Drug Facts

Active ingredient (in each caplet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves:

■ travelers' diarrhea

■ diarrhea

■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products.

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days.
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions

- swallow caplets with water, do not chew
- **adults and children 12 years and over:** 2 caplets (1 dose) every 1/2 hour or 4 caplets (2 doses) every hour as needed for diarrhea
- 2 caplets (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other Information

- **each caplet contains:** salicylate 102 mg
- store in a dry place at 15° - 30° C (59° - 86° F)
- **TAMPER EVIDENT: do not use if printed seal under cap is torn or missing**

croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, silica, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin.

Questions or comments?

1-800-231-4670

RETAIN THIS CARTON FOR FULL LABELING INFORMATION

*CVS™ is not affiliated with the owner of the registered trademark Pepto-Bismol®.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2026 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

V-30287

WARNING: THIS PRODUCT CAN EXPOSE YOU TO CHEMICALS INCLUDING LEAD, WHICH IS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER AND BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. FOR MORE INFORMATION GO TO WWW.P65WARNINGS.CA.GOV.

CVS™

NDC 69842-346-28

STOMACH RELIEF

Bismuth Subsalicylate, **262 mg**

Upset Stomach Reliever/Antidiarrheal

Compare to Pepto-Bismol® active ingredient*

Relieves 5 symptoms:

Heartburn, Indigestion, Upset stomach, Nausea, Diarrhea

40 CAPLETS

1.8120

1.0000

R53825

Drug Facts (continued)

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children**. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions ■ swallow caplets with water, do not chew ■ adults and children 12 years and over: 2 caplets (1 dose) every 1/2 hour or 4 caplets (2 doses) every hour as needed for diarrhea ■ 2 caplets (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)

Drug Facts

Active ingredient (in each caplet)
Bismuth subsalicylate 262 mg

Uses/relieves: ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products.

*CVS™ is not affiliated with the owner of the registered trademark Pepto-Bismol®.



Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2026 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-30287

100% money back guaranteed.
CVS.com/returnpolicy



NDC 69842-346-28

STOMACH RELIEF

Bismuth Subsaliicylate, 262 mg
Upset Stomach Reliever / Antidiarrheal

Compare to Pepto-Bismol® active ingredient*

40
CAPLETS

Relieves 5 Symptoms:
Heartburn, Indigestion,
Upset stomach,
Nausea, Diarrhea



actual size

RETAIN THIS CARTON FOR FULL LABELING INFORMATION

#688802



Drug Facts (continued)

■ do not exceed 8 doses (16 caplets) in 24 hours ■ use until diarrhea stops but not more than 2 days

■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each caplet contains: salicylate 102 mg ■ Store in a dry place at 15° - 30°C (59° - 86°F) ■ TAMPER EVIDENT: do not use if printed seal under cap is torn or missing

Inactive ingredients croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polydioxane, silica, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin.

Questions or comments? 1-800-231-4670

WARNING: THIS PRODUCT CAN EXPOSE YOU TO CHEMICALS INCLUDING LEAD, WHICH IS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER AND BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. FOR MORE INFORMATION GO TO WWW.P66WARRNING.CA.GOV.

BISMUTH SUBSALICYLATE
bismuth subsalicylate tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-346
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TRIACETIN (UNII: XHX3C3X673)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	V
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-346-28	1 in 1 CARTON	07/06/2018	
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69842-346-12	1 in 1 CARTON	07/06/2018	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	07/06/2018	

Labeler - CVS Pharmacy, Inc. (062312574)

Revised: 2/2026

CVS Pharmacy, Inc.