

**ACETAMINOPHEN, ASPIRIN, CAFFEINE- acetaminophen, aspirin,
caffeine tablet, film coated
CVS Pharmacy, Inc.**

5004-CVS

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses

■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ a cold ■ arthritis ■ toothache ■ premenstrual and menstrual cramps

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ with other drugs containing acetaminophen ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of **caffeine**-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen
- any new symptoms appear.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- **adults and children 12 years and over:**
- take 2 caplets every 6 hours
- do not take more than 8 caplets in 24 hours
- **children under 12 years:** ask a doctor

Other Information

- Store in a dry place at 15° - 30°C (59° - 86°F)
- use by expiration date on package

hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, starch, stearic acid, talc, titanium dioxide, triglycerides

Questions or comments?

1-800-231-4670

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by HALEON, owner of the registered trademark Excedrin®.

**Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2024 CVS/pharmacy
CVS.com® 1-800-SHOP CVS**

Value Size 300 Caplets

CVSHealth

Compare to the active ingredients in Excedrin® Extra Strength Caplets†

EXTRA STRENGTH Headache Relief

ACETAMINOPHEN, ASPIRIN (NSAID)* & CAFFEINE

Pain reliever ; Pain reliever aid

300 Caplets

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

VALUE SIZE 300 CAPLETS

CVS Health. Compare to the active ingredients in Excedrin® Extra Strength Caplets†

EXTRA STRENGTH Headache Relief

ACETAMINOPHEN, ASPIRIN (NSAID)* & CAFFEINE

Pain reliever;
Pain reliever aid



Actual Size

300 CAPLETS

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
<p>Active ingredients (in each caplet)</p> <p>Acetaminophen 250 mg Aspirin 250 mg (NSAID) Caffeine 65 mg</p> <p>†nonsteroidal anti-inflammatory drug</p>
<p>Uses</p> <p>temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ a cold ■ arthritis ■ toothache ■ menstrual and menstrual cramps</p>
<p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 capsules in 24 hours, which is the maximum daily amount ■ 3 or more alcoholic drinks every day while using this product</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product</p>

#966960



0 50428 31133 2

CVS Quality Money Back Guarantee

RECEIVE FEEL FREE

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02896
©2024, CVS/pharmacy
CVS.com® 1-800-SHOP-CVS
V-18611

7.0"

INSIDE OF LABEL

Drug Facts (continued)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ an allergic reaction occurs. Seek medical help right away ■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ painful area is red or swollen ■ any new symptoms appear. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a poison control center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years of age and over:
 - take 2 caplets every 6 hours
 - do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- Store in a dry place at 15° – 30°C (59° – 86°F) ■ use by expiration date on package

Inactive ingredients hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, starch, stearic acid, talc, titanium dioxide, triglycerides

Questions or comments? 1-800-231-4670

This product is not manufactured or distributed by HALEOM, owner of the registered trademark Excedrin®.

NO COPY
ADHESIVE ZONE

ACETAMINOPHEN, ASPIRIN, CAFFEINE

acetaminophen, aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-541
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	2793
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-541-06	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/22/2018	
2	NDC:69842-541-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	06/22/2018	
3	NDC:69842-541-01	1 in 1 CARTON	11/03/2020	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:69842-541-16	1 in 1 CARTON	04/06/2021	
4		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M013

06/22/2018

Labeler - CVS Pharmacy, Inc. (062312574)

Revised: 5/2024

CVS Pharmacy, Inc.