

SODIUM FLUORIDE 1.1%- sodium fluoride gel, dentifrice
Burel Pharmaceuticals LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

SODIUM FLUORIDE 1.1%
DENTAL GEL

SODIUM FLUORIDE 1.1% Prescription Dental Gel

DESCRIPTION:

Self-topical neutral fluoride gel containing 1.1% (w/v) sodium fluoride for use as a dental caries preventative in pediatric patients and adults. This prescription is not a dentifrice.

Active Ingredient:

Sodium Fluoride 1.1% (w/v)

Inactive Ingredients:

water (purified), sorbitol, hydroxyethylcellulose, flavor, poloxamer 407, sodium saccharin, methylparaben, propylparaben, FD&C Blue No.1.

CLINICAL PHARMACOLOGY:

Frequent topical applications to the teeth with preparations having a relatively high fluoride content increase tooth resistance to acid dissolution and enhance penetration of the fluoride ion into tooth enamel.

INDICATIONS AND USAGE:

A dental caries preventive; for once daily self-applied topical use. It is well established that 1.1% sodium fluoride is safe and extraordinarily effective as a caries preventive when applied frequently with mouthpiece applicators. ¹⁻⁴ Sodium Fluoride 1.1% Dental Gel in a squeeze-tube is easily applied onto a toothbrush. This prescription dental gel should be used once daily following use of your regular toothpaste unless otherwise instructed by your dental professional. May be used whether or not drinking water is fluoridated, since topical fluoride cannot produce fluorosis. (See WARNINGS for exception.)

CONTRAINDICATIONS:

Do not use in pediatric patients under age 6 years unless recommended by a dentist or

physician.

WARNINGS:

Prolonged daily ingestion may result in various degrees of dental fluorosis in pediatric patients under age 6 years, especially if the water fluoridation exceeds 0.6 ppm, since younger pediatric patients frequently can not perform the brushing process without significant swallowing. Use in pediatric patients under age 6 years requires special supervision to prevent repeated swallowing of dental gel, which could cause dental fluorosis. Read directions carefully before using.

Keep out of reach of infants and children.

PRECAUTIONS:

Not for systemic treatment. **DO NOT SWALLOW.**

Carcinogenesis, Mutagenesis, Impairment of Fertility:

In a study conducted in rodents, no carcinogenesis was found in male and female mice and female rats treated with fluoride at dose levels ranging from 4.1 to 9.1 mg/kg of body weight. Equivocal evidence of carcinogenesis was reported in male rats treated with 2.5 and 4.1 mg/kg of body weight. In a second study, no carcinogenesis was observed in rats, males or females, treated with fluoride up to 11.3 mg/kg of body weight. Epidemiological data provide no credible evidence for an association between fluoride, either naturally occurring or added to drinking water, and risk of human cancer. Fluoride ion is not mutagenic in standard bacterial systems. It has been shown that fluoride ion has potential to induce chromosome aberrations in cultured human and rodent cells at doses much higher than those to which humans are exposed. *In vivo* data are conflicting. Some studies report chromosome damage in rodents, while other studies using similar protocols report negative results. Potential adverse reproductive effects of fluoride exposure in humans have not been adequately evaluated. Adverse effects on reproduction were reported for rats, mice, fox, and cattle exposed to 100 ppm or greater concentrations of fluoride in their diet or drinking water. Other studies conducted in rats demonstrated that lower concentrations of fluoride (5 mg/kg of body weight) did not result in impaired fertility and reproductive capabilities.

Pregnancy:

Pregnancy Category B. It has been shown that fluoride crosses the placenta of rats, but only 0.01% of the amount administered is incorporated in fetal tissue. Animal studies (rats, mice, rabbits) have shown that fluoride is not a teratogen. Maternal exposure to 12.2 mg fluoride/kg of body weight (rats) or 13.1 mg/kg of body weight (rabbits) did not affect the litter size or fetal weight and did not increase the frequency of skeletal or visceral malformations. There are no adequate and well-controlled studies in pregnant women. However, epidemiological studies conducted in areas with high levels of naturally fluoridated water showed no increase in birth defects. Heavy exposure to fluoride during in utero development may result in skeletal fluorosis, which becomes evident in childhood.

Nursing Mothers:

It is not known if fluoride is excreted in human milk. However, many drugs are excreted in milk, and caution should be exercised when products containing fluoride are administered to a nursing woman. Reduced milk production was reported in farm-raised fox when the animals were fed a diet containing a high concentration of fluoride (98-137 mg/kg of body weight). No adverse effects on parturition, lactation, or offspring were seen in rats administered fluoride up to 5 mg/kg of body weight.

Pediatric Use:

The use of Sodium Fluoride 1.1% Dental Gel in pediatric age groups 6 to 16 years as a caries preventive is supported by pioneering clinical studies with 1.1% sodium fluoride gels in mouth trays in students age 11-14 years conducted by Englander, et al.²⁻⁴ Safety and effectiveness in pediatric patients below the age of 6 years have not been established. Please refer to the CONTRAINDICATIONS and WARNINGS sections.

ADVERSE REACTIONS:

Allergic reactions and other idiosyncrasies have been rarely reported. To report SUSPECTED ADVERSE REACTIONS, contact Sheffield Pharmaceuticals, LLC. at 1-800-222-1087 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE:

Accidental ingestion of large amounts of fluoride may result in acute burning in the mouth and sore tongue. Nausea, vomiting, and diarrhea may occur soon after ingestion (within 30 minutes) and are accompanied by salivation, hematemesis, and epigastric cramping abdominal pain. These symptoms may persist for 24 hours. If less than 5 mg fluoride/kg body weight (i.e., less than 2.3 mg fluoride/lb body weight) have been ingested, give calcium (e.g., milk) orally to relieve gastrointestinal symptoms and observe for a few hours. If more than 5 mg fluoride/kg body weight (i.e., more than 2.3 mg fluoride/lb body weight) have been ingested, induce vomiting, give orally soluble calcium (e.g., milk, 5% calcium gluconate or calcium lactate solution) and immediately seek medical assistance. For accidental ingestion of more than 15 mg fluoride/kg of body weight (i.e., more than 6.9 mg fluoride/lb body weight), induce vomiting and admit immediately to a hospital facility. A treatment dose (a thin ribbon) of Sodium Fluoride 1.1% Dental Gel contains 2 mg fluoride. A 2 oz. tube contains 255 mg fluoride.

DOSAGE AND ADMINISTRATION:

Follow these instructions unless otherwise instructed by your dental professional:

1. Adults and pediatric patients 6 years of age or older, apply a thin ribbon of gel to the teeth with a toothbrush or mouth trays once daily for at least one minute, preferably at bedtime.
2. After use, adults expectorate gel. For best results, do not eat, drink, or rinse for 30 minutes. Pediatric patients, age 6-16, expectorate after use and rinse mouth thoroughly.

HOW SUPPLIED:

Fresh Mint

2 oz. (56g) net wt. tube

NDC# 35573-435-56

STORAGE:

Store at Controlled Room Temperature, 20-25°C (68-77°F).

REFERENCES:

1. American Dental Association, Council on Dental Therapeutics, Fluoride compounds, In:Accepted Dental Therapeutics, Ed. 40, Chicago, ADA, 405-407 (1984).
2. H.R. Englander et al., Clinical Anticaries Effects of Repeated Topical Sodium Fluoride Application by Mouthpieces, JADA, 75, 638-644 (1967).
3. H.R. Englander et al., Residual Anticaries Effects of Repeated Topical Sodium Fluoride Application by Mouthpieces, JADA, 78, 783-787 (1969).
4. H.R. Englander et al., Incremental Rates of Dental Caries After Repeated Topical Sodium Fluoride Applications in Children with Lifelong Consumption of Fluoridated Water, JADA, 82, 354-358 (1971).

PRINCIPAL DISPLAY PANEL - 56 g Tube

burelpharma

NDC 35573-435-56

Rx only

Sodium Fluoride 1.1%

Dental Gel

Prescription Brush-on Gel (non-abrasive formula)

FRESH MINT

NET WT. 2.0 OZ. (56 g)



NDC 35573-435-56
Rx Only

Sodium Fluoride 1.1% Dental Gel

FRESH MINT

Prescription Brush-on Gel (non-abrasive formula)

NET WT. 2 OZ. (56 g)

Description: Self-topical neutral 1.1% sodium fluoride for use as a dental caries preventive. This is not a dentifrice.

Directions for daily use:

1. After brushing with toothpaste, rinse as usual. Adults and pediatric patients 6 years of age or older, apply a thin ribbon of gel on the teeth with a toothbrush or mouthtrays for at least one minute, preferably at bedtime.
2. After use, adults expectorate gel. For best results, do not eat, drink or rinse for 30 minutes. Pediatric patients, age 6-16, expectorate gel after use and rinse mouth thoroughly.

Precautions: DO NOT SWALLOW. As with all medications, keep out of reach of infants and children.

Store at controlled room temperature, 20°-25°C (68°-77°F).

See package insert for additional information.

Manufactured for:
Burel Pharmaceuticals, LLC
Mason, OH 45040 USA

Rev. 06/21



2502512



3 35573 43556 1

PRINCIPAL DISPLAY PANEL - 56 g Carton

burelpharma NDC 35573-435-56



Rx only


Sodium Fluoride 1.1%

Dental Gel

Prescription Brush-on Gel (non-abrasive formula)

FRESH MINT

NET WT 2.0 oz (56g)



SODIUM FLUORIDE 1.1%

sodium fluoride gel, dentifrice

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:35573-435
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU4080)	SODIUM FLUORIDE	5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)	
POLOXAMER 407 (UNII: TUF21VW3M2)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35573-435-56	1 in 1 CARTON	02/28/2020	
1		56 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		02/28/2020	

Labeler - Burel Pharmaceuticals LLC (609436204)

Revised: 11/2025

Burel Pharmaceuticals LLC