

COLD AND FLU SEVERE, DAY, NIGHT- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr. guaifenesin, phenylephrine hcl
L.N.K. International, Inc.

Quality Plus 44-503A473C-08

Active ingredients (in each caplet) (Daytime Cold & Flu Severe)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - headache
 - sore throat
 - nasal congestion
 - minor aches and pains
 - sinus congestion and pressure
 - sneezing and runny nose (***Nighttime only***)
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (***Daytime only***)

- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- high blood pressure
- liver disease
- diabetes
- thyroid disease
- heart disease
- glaucoma (**Nighttime only**)
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)
- use caution when driving a motor vehicle or operating machinery (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole – do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

**QUALITY
+PLUS**

NDC 50844-547-08

*Compare to active ingredients in
Tylenol® COLD + FLU
SEVERE DAY & NIGHT

**SEVERE
COLD & FLU**

DAY

Acetaminophen,
Dextromethorphan HBr,
Guaifenesin,
Phenylephrine HCl

PAIN RELIEVER/
FEVER REDUCER,
COUGH SUPPRESSANT,
EXPECTORANT,
NASAL DECONGESTANT

ACTUAL
SIZE

**16
Caplets**

NIGHT

Acetaminophen,
Chlorpheniramine maleate,
Dextromethorphan HBr,
Phenylephrine HCl

PAIN RELIEVER/
FEVER REDUCER,
ANTIHISTAMINE,
COUGH SUPPRESSANT,
NASAL DECONGESTANT

ACTUAL
SIZE

8
Caplets

24 Total Caplets

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® COLD + FLU SEVERE DAY & NIGHT.

50844 REV0922A50347308

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

**Do Not Take Daytime and
Nighttime Products at the
Same Time.**

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

QUALITY PLUS
SEVERE
COLD & FLU
DAY & NIGHT

QUALITY PLUS NDC 50844-547-08
SEVERE
COLD & FLU

DAY
Acetaminophen,
Dextromethorphan HBr,
Guaifenesin,
Phenylephrine HCl
PAIN RELIEVER/
FEVER REDUCER,
COUGH SUPPRESSANT,
EXPECTORANT,
NASAL DECONGESTANT

NIGHT
Acetaminophen,
Chlorpheniramine maleate,
Dextromethorphan HBr,
Phenylephrine HCl
PAIN RELIEVER/
FEVER REDUCER,
ANTIHISTAMINE,
COUGH SUPPRESSANT,
NASAL DECONGESTANT

ACTUAL SIZE
16 Caplets

ACTUAL SIZE
8 Caplets

24 Total Caplets



No print/No varnish area
Lot no/Exp date

QUALITY PLUS
SEVERE
COLD & FLU
DAY & NIGHT

B-1603-503A473C-08
REV0922A50347308

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

<p>Drug Facts (continued)</p> <p>Active ingredients (in each caplet) (Daytime Cold & Flu Severe)</p> <ul style="list-style-type: none"> Acetaminophen 325 mg Dextromethorphan HBr 10 mg Guaifenesin 200 mg Phenylephrine HCl 3 mg <p>Active ingredients (in each caplet) (Nighttime Cold & Flu Severe)</p> <ul style="list-style-type: none"> Acetaminophen 325 mg Chlorpheniramine maleate 2 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 3 mg <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> cough headache sore throat nasal congestion sinus congestion and pressure steezing and runny nose (Nighttime only) 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive (Daytime only) helps clear nasal passages relieves cough to help you sleep temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin redness hives itching swelling of the face, lips, tongue, or throat skin rash </p> <p>See throat warning. If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or over-the-counter)
--	--

TYPED LABEL

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

PARENTS:
Learn about how medicine abuse affects your child.
www.StopMedicineAbuse.org

Do Not Take Daytime and Nighttime Products at the Same Time.

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® COLD & FLU SEVERE DAY & NIGHT.
50844 REV0922A50347308
Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

Drug Facts (continued)

nonprescription. If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- high blood pressure
- liver disease
- heart disease
- glaucoma (Nighttime only)
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis (Nighttime only)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Drug Facts (continued)

- taking sedatives or tranquilizers (Nighttime only)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- avoid alcoholic beverages (Nighttime only)
- use caution when driving a motor vehicle or operating machinery (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- each caplet contains: sodium 3 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients (Daytime only)

corn starch, croscopolvidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, croscopolvidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-426-9391

50844
REV0922A50347308
R-1603-503A473C08-DF

Quality Plus 44-503A473C

COLD AND FLU SEVERE, DAY, NIGHT
acetaminophen, chlorpheniramine maleate, dextromethorphan hbr. guaifenesin, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-547
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-547-08	1 in 1 CARTON; Type 0: Not a Combination Product	07/15/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	1 BLISTER PACK	8

Part 1 of 2

COLD AND FLU SEVERE DAY

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:50844-980
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MENTHOL	Imprint Code	44;503
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2021	

Part 2 of 2

COLD AND FLU SEVERE NIGHT

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Item Code (Source)	NDC:50844-981
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -	PHENYLEPHRINE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor	MENTHOL	Imprint Code	44;473
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2021	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-547) , pack(50844-547)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-547)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-547)

Revised: 5/2025

L.N.K. International, Inc.