LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH- acetaminophen tablet Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Lil' Drug Store® Pain Reliever Extra Strength

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 6 tablets (3,000 mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 tablets every 6 hours while symptoms last do not take more than 6 tablets in 24 hours unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- do not use if vial is open or damaged or safety seal has been removed
- store at room temperature 59-86°F (15-30°C)

Inactive ingredients

corn starch, hypromellose, maltodextrin¹, microcrystalline cellulose¹, polyethylene glycol, povidone¹, pregelatinized starch¹, sodium starch glycolate¹, stearic acid, titanium dioxide¹

Questions or comments?

call toll-free 1-877-507-6516 (M-F 8 AM-4:30PM CST)

QUALITY GUARANTEED

Compare to the Active Ingredient in Tylenol® Extra Strength*

Extra Strength Pain Reliever

Acetaminophen, 500 mg Pain Reliever/Fever Reducer

10

Tablets

Lil'

 $DrugStore^{\circledR}$

OPEN HERE FOR DRUG FACTS INFORMATION

Lil' Drug Store Products, Inc. does not own the Tylenol® Extra Strength trademark. McNeil Consumer Healthcare Division of McNeil-PPC, Inc. *This product is not manufactured or distributed by

> Cedar Rapids, IA 52404 9300 Earhart Lane SW Product manufactured for: Lif Drug Store Products, Inc.

(M-F 8 AM-4:30PM CST) call toll-free 1-877-507-6516 Questions or comments?

starch glycolate*, stearic acid, titanium dioxide* *may contain povidone*, pregelatinized starch*, sodium microcrystalline cellulose*, polyethylene glycol, corn starch, hypromellose, maltodextrin*, inactive ingredients

 store at room temperature 59-86°F (15-30°C) safety seal has been removed · do not use it vial is open or damaged or Other information

	Othor into
sak a doctor	children under 12 years
• take 2 tablets every 6 hours while symptoms last of on oit take more than 6 tablets in 24 hours unless directed by a doctor on not use for more than 10 days unless than 10 days unless directed by a doctor	adults and children 12 years and over

adults and (see overdose warning) · do not take more than directed Directions

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose) may cause help or confact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)



Compare to the Active Ingredient in Tylenol® Extra Strength*

Extra Strength
Pain Reliever

Acetaminophen, 500 mg Pain Reliever/Fever Reducer

Tablets



Drug Facts Active ingredient (in each tablet)

Purpose

Acetaminophen 500 mg..Pain reliever/fever reduct

Uses

SAFETY SEALED

- · temporarily relieves minor aches and pains due to:
- · the common cold
- headache · minor pain of arthritis
- · backache · toothache
- · muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings
Liver warning: This product contains
acetaminophen. The maximum daily dose of this product is 6 tablets (3,000 mg) in 24 hours.

- Severe liver damage may occur if you take

 more than 4,000 mg of acetaminophen in 24
- with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

- with any other drug containing acetaminophen Do not use (prescription or nonprescription). If you are not sure whether a drug contains
- acetaminophen, ask a doctor or pharmacist. if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- · fever gets worse or lasts more than 3 days
- new symptoms occur
- · redness or swelling is present

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children.





LIL DRUG STORE PAIN RELIEVER EXTRA STRENGTH

acetaminophen tablet

Droc	luct	Info	rmation
PIUU)

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:66715-9737

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	500 mg

Inactive Ingredients			
Ingredient Name	Strength		
Starch, Corn (UNII: O8232NY3SJ)			
Hypromellose, Unspecified (UNII: 3NXW29V3WO)			
Maltodextrin (UNII: 7CVR7L4A2D)			
Microcrystalline Cellulose (UNII: OP1R32D61U)			
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)			
Povidone, Unspecified (UNII: FZ989GH94E)			
Sodium Starch Glycolate Type A Corn (UNII: AG9B65PV6B)			
Stearic Acid (UNII: 4ELV7Z65AP)			
Titanium Dioxide (UNII: 15FIX9 V2JP)			
Sodium Starch Glycolate Type A Potato (UNII: 5856J3G2A2)			

Product Characteristics			
Color	WHITE	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	FR33
Contains			

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:66715-9737-3	3 in 1 CARTON	0 1/27/20 11	08/30/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product			
2	NDC:66715-9737-4	10 in 1 VIAL; Type 0: Not a Combination Product	0 1/27/20 11		

Marketine	Tnfaun	nation
Marketing	g muori	nauon

Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End Date

OTC MONOGRAPH NOT FINAL	part343	0 1/27/20 11	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 8/2019 Lil' Drug Store Products, Inc.