

**BISACODYL- bisacodyl suppositories suppository**  
**Oncor Pharmaceuticals**

-----  
**Bisacodyl Suppositories**

<b>Active ingredient (in each suppository)</b>	<b>Purpose</b>
Bisacodyl USP 10 mg.....	Stimulant Laxative

**Uses**

- relieves occasional constipation
- this product generally produces a bowel movement in 15 minutes 1 hour

**Warning**

**for rectal use only**

**Do not use** when abdominal pain, nausea, or vomiting are present.

**Ask a doctor before use if you have** a sudden change in bowel habits that lasts over a period of 2 weeks.

**When using this product** you may experience

- rectal burning
- mild cramps
- abdominal discomfort
- faintness

**Stop use and ask a doctor if**

- you have rectal bleeding or you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.
- you need to use a laxative for more than one week

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

You may report side effects by calling Oncor Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.

**Directions**

- tear off one suppository from the strip
- peel apart the wrapper
- continue peeling apart to almost the full length of the suppository

- gently remove the suppository from the wrapper
- lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out
- retain it for 15 to 20 minutes
- **Do not exceed recommended dose**
- if suppository seems soft, place in refrigerator for a short time before use.

---

Adults and children 12 years and over	1 suppository once daily
Children 6 to under 12 years	1/2 suppository once daily
Children under 6 years	Ask a doctor

---

### **Other information**

store at temperatures below 25 °C (77 °F)

### **Inactive ingredient**

hydrogenated vegetable oil

**TAMPER EVIDENT:**SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED.

### **PRINCIPAL DISPLAY PANEL**

Oncor Pharmaceuticals

**NDC 83720-550-25**

12 Suppositories

10 mg each

Bisacodyl Suppositories

Fast Acting Stimulant Laxative

Manufactured In USA For :

Oncor Pharmaceuticals  
6755 Business Pkwy Suite 202  
Elkridge, MD 21075  
United States.

Rev. 12/24

**NCOR**  
 PHARMACEUTICALS  
 NDC 83720-550-25

12 Suppositories  
 10 mg each

# Bisacodyl Suppositories

Fast Acting Stimulant Laxative


TAMPER EVIDENT: SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED.

Bisacodyl  
 Suppositories  
 Fast Acting Stimulant Laxative

## Bisacodyl Suppositories

Fast Acting Stimulant Laxative

Bisacodyl  
 Suppositories  
 Fast Acting Stimulant Laxative

<p><b>Drug Facts</b>  <b>Active ingredient (in each suppository)</b>          Bisacodyl USP 10 mg</p>	<p><b>Purpose</b>          Stimulant Laxative</p>							
<p><b>Uses</b>          • relieves occasional constipation          • this product generally produces a bowel movement in 15 minutes to 1 hour</p>	<p><b>Directions</b>          • tear off one suppository from the strip • peel apart the wrapper          • continue peeling apart to almost the full length of the suppository (see illustration)          • gently remove the suppository from the wrapper          • lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out.          • retain it for 15 to 20 minutes • Do not exceed recommended dose          • if suppository seems soft, place in refrigerator for a short time before use.</p>							
<p><b>Warning for rectal use only</b>          Do not use when abdominal pain, nausea, or vomiting are present.</p>	<p><b>When using this product</b> you may experience          • rectal burning • mild cramps • abdominal discomfort • faintness</p>	<table border="1"> <tr> <td>Adults and children 12 years and over</td> <td>1 suppository once daily</td> </tr> <tr> <td>Children 6 to under 12 years</td> <td>1/2 suppository once daily</td> </tr> <tr> <td>Children under 6 years</td> <td>Ask a doctor</td> </tr> </table>	Adults and children 12 years and over	1 suppository once daily	Children 6 to under 12 years	1/2 suppository once daily	Children under 6 years	Ask a doctor
Adults and children 12 years and over	1 suppository once daily							
Children 6 to under 12 years	1/2 suppository once daily							
Children under 6 years	Ask a doctor							
<p><b>Ask a doctor before use if you have</b> a sudden change in bowel habits that lasts over a period of 2 weeks.</p>	<p><b>Stop use and ask a doctor if</b>          • you have rectal bleeding or you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.          • you need to use a laxative for more than one week</p>	<p>You may report side effects by calling Oncor Pharmaceuticals (9 a.m. to 5 p.m. EST), at 1-443-876-7600 or FDA at 1-800-FDA-1088.</p>						
<p><b>If pregnant or breast-feeding</b>, ask a health professional before use.  <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p><b>Other information</b>          • store at temperatures below 25°C (77°F)</p>	<p><b>Inactive ingredient</b> hydrogenated vegetable oil</p>						

Manufactured In USA For :  
 Oncor Pharmaceuticals  
 6755 Business Pkwy Suite 202  
 Elkridge, MD 21075  
 United States.



Rev. 12/24

## BISACODYL

bisacodyl suppositories suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83720-550
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

## Inactive Ingredients

Ingredient Name	Strength
PALM KERNEL OIL (UNII: B0S90M0233)	

## Product Characteristics

Color	white	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83720-550-25	12 in 1 CARTON; Type 0: Not a Combination Product	02/01/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/01/2025	

**Labeler** - Oncor Pharmaceuticals (119032580)

**Registrant** - Oncor Pharmaceuticals (119032580)