

**MAXIMUM STRENGTH MUCINEX FAST-MAX COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution**  
**RB Health (US) LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Maximum Strength Mucinex® Fast-Max® Cold & Flu**

**Drug Facts**

<b>Active ingredients (in each 20 mL)</b>	<b>Purposes</b>
<b>Acetaminophen 650 mg</b>	<b>Pain reliever/fever reducer</b>
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - stuffy nose
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### **Other information**

- each 20 mL contains: **sodium 10 mg**
- store between 20-25°C (68-77°F)
- do not refrigerate

### **Inactive ingredients**

anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

### **Questions?**

**1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser  
Parsippany, NJ 07054-0224

### **PRINCIPAL DISPLAY PANEL - 180 mL Bottle Label**

#### **MAXIMUM STRENGTH**

NDC 63824-527-03

**Mucinex®**  
**FAST-MAX®**

#### **COLD & FLU**

Acetaminophen – Pain Reliever/Fever Reducer  
Dextromethorphan HBr – Cough Suppressant  
Guaifenesin – Expectorant

Phenylephrine HCl - Nasal Decongestant

HEADACHE

BODY PAIN

SORE THROAT

FEVER

CHEST CONGESTION

COUGH

**ALL IN**

**ONE\***

NASAL CONGESTION

SINUS CONGESTION

SINUS PRESSURE

*ORANGE & PINEAPPLE FLAVOR*

**6 FL OZ (180 mL)**

**FOR AGES 12+**

MAXIMUM STRENGTH

NDC 63824-527-03



### COLD & FLU

Acetaminophen – Pain Reliever/Fever Reducer  
Dextromethorphan HBr – Cough Suppressant  
Guaifenesin – Expectorant  
Phenylephrine HCl – Nasal Decongestant



ORANGE & PINEAPPLE FLAVOR

3064040

6 FL OZ (180 mL) FOR AGES 12+

030218

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Maximum Strength per 4-hour dose  
Tamper evident: Do not use if neckband on bottle cap is broken or missing.

\*Helps to relieve these symptoms day or night

### PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Dist. by: Beckitt Benckiser  
Parsippany, NJ 07054-0224  
© 2018 RB  
020618

LOT: 3064041  
EXP:  
MADE IN:

### Drug Facts

Active ingredients (in each 20 mL) Purposes

Acetaminophen 650 mg...Pain reliever/fever reducer  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Guaifenesin 400 mg.....Expectorant  
Phenylephrine HCl 10 mg.....Nasal decongestant

### Uses

- temporarily relieves these common cold and flu symptoms:
  - cough
  - minor aches and pains
  - headache
  - sinus congestion and pressure
  - nasal congestion
  - sore throat
  - stuffy nose
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 6 doses in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks daily while using this product

### Drug Facts (continued)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin  
When using this product do not use



If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure

PEEL HERE

**more than directed**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see **Overdose warning**)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided

**Drug Facts (continued)**

- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients** anhydrous citric acid, D&C yellow no. 10, edetate disodium, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions?**  
1-866-MUCINEX (1-866-682-4639)  
You may also report side effects to this phone number.

  
 HEALTH • HYGIENE • HOME  
 Please visit our website  
[www.mucinex.com](http://www.mucinex.com)



## MAXIMUM STRENGTH MUCINEX FAST-MAX COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:63824-527

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-527-03	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	07/28/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/28/2018	

**Labeler** - RB Health (US) LLC (081049410)

## Establishment

Name	Address	ID/FEI	Business Operations
Reckitt Benckiser (UK) Ltd.		227363660	manufacture(63824-527)

## Establishment

Name	Address	ID/FEI	Business Operations
RB Salute Mexico, S.A. de C.V.		812814950	manufacture(63824-527)

Revised: 11/2022

RB Health (US) LLC