

**ADULT LOW DOSE ASPIRIN- aspirin tablet, delayed release**  
**VALU MERCHANDISERS COMPANY**

-----  
**1146-BST-2024-0613**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID\*)

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- temporarily relieves minor aches and pains
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- diabetes
- gout
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### **Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

## **Inactive ingredients**

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, triethyl citrate

## **Questions or comments?**

1-844-292-1112

## **PRINCIPAL DISPLAY PANEL**

Best Choice®

Compare to the active ingredient in St.Joseph® Low Dose Aspirin†

LOW DOSE

Aspirin 81 mg

PAIN RELIEVER (NSAID)

Aspirin Regimen\*\*

Safety Coated

Talk to your doctor or other healthcare provider before using this product for your heart

Actual Size

81mg each

100 ENTERIC COATED TABLETS



**ADULT LOW DOSE ASPIRIN**  
 aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-246
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

## Product Characteristics

Color	orange (PEACH)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-246-03	1 in 1 CARTON	05/01/2024	06/30/2028
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2024	06/30/2028

Revised: 12/2025

VALU MERCHANDISERS COMPANY