

RID LICE TREATMENT COMPLETE KIT- piperonyl butoxide, pyrethrum extract
Oystershell Consumer Health Inc

RID® Lice Treatment Complete Kit

Drug Facts

Active ingredients

Piperonyl butoxide (4%)

Pyrethrum extract (equivalent to 0.33% pyrethrins)

Purpose

Lice treatment

Uses

treats head, pubic (crab), and body lice

Warnings

For external use only

Do not use

- near the eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.
- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you are

allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

● ● Important: Read warnings before use

Adults and children 2 years and over:

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

Treat

- apply thoroughly to **DRY HAIR** or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for 10 minutes, but no longer
- use warm water to form a lather, shampoo, then thoroughly rinse
- for head lice, towel dry hair and comb out tangles

Remove lice and their eggs (nits)

- use the RID® Lice Comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments

Children under 2 years: ask a doctor

Other information

- keep carton for important product information
- see Consumer Information Insert for additional information
- avoid excessive heat

Inactive ingredients

ammonium laureth sulfate, dehydrated alcohol, fragrance, isopropyl alcohol, PEG-25 hydrogenated castor oil, polyquaternium-10, purified water

Questions or comments?

1-800-RID-LICE
(1-800-743-5423)

Principal Display Panel

RID®

**LICE TREATMENT
COMPLETE KIT**
LICE TREATMENT

Easy-to-Follow 3-Step Process—
Instructions Inside

- ① **TREAT**
- ② **REMOVE**
- ③ **CONTROL**

**RID
SUPPORT**

CALL AN EXPERT
1-800-RID-LICE

HOW-TO VIDEOS

TRATAMIENTO DE PIOJOS—KIT COMPLETO
Se incluyen instrucciones en español

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautions

SHAMPOO NET WT. 4 FL OZ (118 mL)
COMB-OUT SPRAY NET WT. 2 FL OZ (59 mL)
RID® HOME LICE, BEDBUG & DUST MITE SPRAY
NET WT. 3 OZ (85.1 g)
1 PATENTED RID® LICE COMB

**RID® provides everything you
need to treat lice in your family
and in your home.**

Step 1 TREAT

- **Apply RID® Lice Killing Shampoo** according to label directions.
- Repeat this step 7 to 10 days later to kill any newly hatched lice.

Step 2 REMOVE

- After Step 1, apply a small amount of RID® Lice & Egg Comb-Out Spray and use a wide-tooth comb to remove any tangles.
- Comb out the eggs and nits in the hair with RID® Lice comb (included), using

additional spray to remove tangles if needed.

Step 3 CONTROL

- Use RID® Home Lice, Bedbug & Dust Mite Spray to kill lice and their eggs on mattresses, furniture, car interiors, and other non-washable items.

IMPORTANT: Read detailed directions inside before using.

This kit contains 4 products (Lice Killing Shampoo, Lice & Egg Comb-Out Spray, Lice Comb and RID® Home Lice, Bedbug & Dust Mite Spray **to treat lice in your family and home. The 4oz. bottle of shampoo and 2oz bottle of comb-out spray should be enough product to treat one person one time.** However, the amount of shampoo needed will vary by hair length and you may need additional shampoo for the second treatment 7 to 10 days later. **See below for guidelines.**

Lice Killing Shampoo

Hair Length	Approximate Amount For 1 Adult/Child:
Short (ear length or shorter)	First Application: 1 oz - 2 oz <u>Second Application: 1 oz - 2 oz</u> Total: 2 oz - 4 oz
Medium (shoulder length)	First Application: 2 oz - 3 oz <u>Second Application: 2 oz - 3 oz</u> Total: 4 oz - 6 oz
Long (past shoulder length)	First Application: 3 oz - 4 oz <u>Second Application: 3 oz - 4 oz</u> Total: 6 oz - 8 oz



LICE TREATMENT COMPLETE KIT

L I C E T R E A T M E N T

Easy-to-Follow 3-Step Process—
Instructions Inside



- 1 TREAT
- 2 REMOVE
- 3 CONTROL

**RID
SUPPORT**

**CALL AN EXPERT
1-800-RID-LICE**

HOW-TO VIDEOS

TRATAMIENTO DE PIOJOS—KIT COMPLETO
Se incluyen instrucciones en español

SHAMPOO NET WT. 4 FL OZ (118 mL)
 COMB-OUT SPRAY NET WT. 2 FL OZ (59 mL)
 RID® HOME LICE, BEDBUG & DUST MITE SPRAY
 NET WT. 3 OZ (85.1 g)
 1 PATENTED RID® LICE COMB

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel for Additional Precautions

RID® provides everything you need to treat lice in your family and in your home.

Step 1 TREAT

- Apply RID® Lice Killing Shampoo according to label directions.
- Repeat this step 7 to 10 days later to kill any newly hatched lice.

Step 2 REMOVE

- After Step 1, apply a small amount of RID® Lice & Egg Comb-Out Spray and use a wide-tooth comb to remove any tangles.
- Comb out the eggs and nits in the hair with RID® Lice Comb (included), using additional spray to remove tangles if needed.

Step 3 CONTROL

- Use RID® Home Lice, Bedbug & Dust Mite Spray to kill lice and their eggs on mattresses, furniture, car interiors, and other non-washable items.

IMPORTANT: Read detailed directions inside before using.

This kit contains 4 products (Lice Killing Shampoo, Lice & Egg Comb-Out Spray, Lice Comb and RID® Home Lice, Bedbug & Dust Mite Spray) to treat lice in your family and home. The 4oz. bottle of shampoo and 2oz bottle of comb-out spray should be enough product to treat one person one time. However, the amount of shampoo needed will vary by hair length and you may need additional shampoo for the second treatment 7 to 10 days later. See below for guidelines.

Lice Killing Shampoo

Hair Length	Approximate Amount For 1 Adult/Child:
Short (ear length or shorter)	First Application: 1 oz - 2 oz
	Second Application: 1 oz - 2 oz
	Total: 2 oz - 4 oz
Medium (shoulder length)	First Application: 2 oz - 3 oz
	Second Application: 2 oz - 3 oz
	Total: 4 oz - 6 oz
Long (past shoulder length)	First Application: 3 oz - 4 oz
	Second Application: 3 oz - 4 oz
	Total: 6 oz - 8 oz

LICE KILLING SHAMPOO

Drug Facts	
Active Ingredients	Purpose
Piperonyl butoxide (4%) Pyrethrum extract (equivalent to 0.33% pyrethrins)	Lice treatment Lice treatment
Uses treats head, pubic (crabs), and body lice	
Warnings	
For external use only	
Do not use	
<ul style="list-style-type: none"> near the eyes inside the nose, mouth, or vagina on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas. If you have ever had an allergic reaction to this product or any of its ingredients 	
Ask a doctor before use if you are allergic to ragweed. May cause breathing difficulty or an asthmatic attack.	
When using this product	
<ul style="list-style-type: none"> keep eyes tightly closed and protect eyes with a washcloth or towel if product gets in eyes, flush with water right away scalp itching or redness may occur 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> breathing difficulty occurs eye irritation occurs skin or scalp irritation continues or infection occurs 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> Important: Read warnings before use Adults and children 2 years and over: Inspect check each household member with a magnifying glass in bright light for lice/nits (eggs) look for tiny nits near scalp, beginning at back of neck and behind ears examine small sections of hair at a time urika dandruff which moves when touched, nits stick to the hair if either lice or nits are found, treat with this product 	
Treat	
<ul style="list-style-type: none"> apply thoroughly to DRY HAIR or other affected area. For head lice, first apply behind ears and to back of neck. allow product to remain for 10 minutes, but no longer use warm water to form a lather, shampoo, then thoroughly rinse for head lice, towel dry hair and comb out tangles 	
Remove lice and their eggs (nits)	
<ul style="list-style-type: none"> use the RID® Lice Comb. Remove any remaining nits by hand (using a throw-away glove). hair should remain slightly damp while removing nits if hair dries during combing, dampen slightly with water for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours. lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp. pin back each strand of hair after combing clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back. after combing, thoroughly recheck for lice/nits. Repeat combing if necessary. check daily for any lice/nits that you missed a second treatment must be done in 7 to 10 days to kill any newly hatched lice if infestation continues, see a doctor for other treatments Children under 2 years: ask a doctor 	

Drug Facts (continued)
Other information
<ul style="list-style-type: none"> keep carton for important product information see Consumer Information Insert for additional information avoid excessive heat
Inactive ingredients ammonium lauryl sulfate, dehydrated alcohol, fragrance, isopropyl alcohol, PEG-25 hydrogenated castor oil, polyquaternium-10, purified water
Questions or comments? 1-800-RID-LICE (1-800-743-5423)

LICE & EGG COMB-OUT SPRAY

Product Information
Use
RID® Lice & Egg Comb-Out Spray detangles hair in order to make lice and egg comb-out easier.
Warnings
For external use only. Keep out of reach of children. Avoid contact with eyes. Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Directions For Use
<ul style="list-style-type: none"> Before using, read Consumer Information Insert for complete directions. Use AFTER shampooing with RID® Lice Killing Shampoo. towel dry hair. Apply a small amount of RID® Lice & Egg Comb-Out Spray and use a wide-tooth comb to remove any tangles. Comb out the dead lice, eggs, and nits, one small section at a time, with the enclosed RID® Lice Comb. (See insert for additional combing instructions). This step is very important. Use additional RID® Lice & Egg Comb-Out Spray to remove tangles, if needed, during the comb-out process. It is important to wash in hot water above 54°C (130°F) all clothing, bedding, towels, and hair products (combs, brushes) used by infested persons in the two days prior to treatment.
Other Information
<ul style="list-style-type: none"> It is important to wash in hot water (130°F) all clothing, bedding, towels, and hair products (combs, brushes) used by infested persons in the two days prior to treatment. Dry clean non-washable fabrics. RID® Lice & Egg Comb-Out Spray does not kill lice or their eggs. Use it is product only after treating with RID® Lice Killing Shampoo. Store in a cool, dry place. Keep from freezing.
Ingredients
Water, Glycerin, Propylene Glycol, Polysorbate 20, Cetrimonium Chloride, Amoldimethicone, Phenylethanol, Fragrance, DL Panthanol, Tris(ortho-12, Methylchloroisothiazolinone, Methylisothiazolinone.
Questions?
1-800-RID-LICE (1-800-743-5423)

RID® HOME LICE, BEDBUG & DUST MITE SPRAY

ACTIVE INGREDIENT:	0.50%
Permethrin	
OTHER INGREDIENTS**:	99.50%
Total	100.00%
*cis/trans ratio: Max. 55% (+/-) cis and Min. 45% (+/-) trans.	
**Contains petroleum distillate	
PRECAUTION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)	
KEEP OUT OF REACH OF CHILDREN	
CAUTION	
FIRST AID: If Swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.	
If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-RID-LICE (1-800-743-5423). NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia.	
PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS & DOMESTIC ANIMALS	
CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.	
PHYSICAL OR CHEMICAL HAZARDS	
Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54°C) may cause bursting.	
DIRECTIONS FOR USE	
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.	
READ ENTIRE LABEL BEFORE EACH USE - Use Restrictions	
<ul style="list-style-type: none"> Do not allow adults, children, or pets to enter until vapors, mists and aerosols have dispersed, and the treated area has been thoroughly ventilated. Do not allow adults, children, or pets to enter the treated area until sprays have dried. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Remove pets, birds and cover any water inhabited by fish (such as aquariums and ornamental fish ponds) before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on. This product is not for use on humans. If lice infestations should occur on humans, use a product labeled for use on humans. 	

<ul style="list-style-type: none"> Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
SHAKE WELL BEFORE USING
Remove protective cap, hold container upright with nozzle away from you. Depress valve and spray from a distance of 8-10 inches (12-15 inches for baseboards, moldings and floors). Test product on an inconspicuous area before applying to check for possible staining or discoloration. Inspect again after drying. Spray to the point of dampness but not run-off. Over application may cause damage. Will not stain surfaces or fabrics where water alone causes no damage.
To Kill Lice and Louse Eggs: Spray entire area to be treated. Spray each square foot for three seconds. Allow all sprayed articles to dry thoroughly before use. Spray only those parts of bedding, including mattresses and furniture that can be either laundered or dry cleaned. Do not use on sheets or pillowcases.
Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without a cover. Repeat if infestation occurs, but not more than once every two weeks.
To Kill House Dust Mites (Dermatophagoides spp.): RID® Home Lice, Bedbug & Dust Mite Spray helps control house dust mites that may cause asthma, hay fever, rhinitis, watery eyes and sneezing. One treatment lasts for up to 8 weeks. On Upholstered Furniture, carpets and curtains: - test fabric in an inconspicuous area to insure eyes will not bleed. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray all fabric surfaces until damp. Do not use on delicate fabrics. Allow furniture to dry thoroughly before using. On Carpets: - spray over entire carpet until damp. Hold the nozzle approximately 12 to 18 inches from the carpet edge and spray in a sweeping motion. Use an overlapping pattern to insure total coverage. After application, brush the surface of the carpet with a broom or brush to insure deep penetration. Allow carpet to dry thoroughly before reentering the treated area. Drying time can be accelerated by using a fan. On Mattresses: - spray the top, bottom and sides of mattresses until damp, paying particular attention to tufts and seams. Allow to dry thoroughly before use. After mattress is dry, cover with mattress cover and sheet. Do not use mattress without a cover. Vacuum entire area once dry. Since vacuuming stirs dust up in the air those with allergies should have another person vacuum or should wear a dust mask and use vacuum with a high efficiency particulate (HEPA) filter.
STORAGE AND DISPOSAL
Pesticide Storage: Store in a cool, dry area away from children and pets and away from heat, sparks and open flame. Do not transport or store below 32°F (0°C).
Pesticide Disposal and Container Handling: Do Not Puncture or Incinerate. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. This container may be recycled in aerosol recycling centers. At present, there are only a few such centers in the country. Before offering for recycling, empty the can by using the product according to the label.
Dist. by: Oystershell Consumer Health, Inc. 409 Lancaster Ave. Wayne, PA 19087 Comb is made in China. Detangling Spray is made in Canada. Shampoo is Made in the USA or US & Imported Materials RID and the RID logo are registered trademarks of Oystershell Consumer Health, Inc. EPA Reg. No. 1021-2685-98143 EPA Est. No. 100939-KS-1 NDC 82392-101-04 Pat. No. 7,909,042

RID LICE TREATMENT COMPLETE KIT

piperonyl butoxide, pyrethrum extract kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82392-101
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82392-101-04	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	01/05/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL

Part 1 of 1

RID LICE KILLING

piperonyl butoxide and pyrethrum extract shampoo

Product Information

Item Code (Source) NDC:82392-100

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PIPERONYL BUTOXIDE (UNII: LWK91TU9AH) (PIPERONYL BUTOXIDE - UNII:LWK91TU9AH)	PIPERONYL BUTOXIDE	40 mg in 1 mL
PYRETHRUM EXTRACT (UNII: ZUM06L90GV) (PYRETHRUM EXTRACT - UNII:ZUM06L90GV)	PYRETHRUM EXTRACT	0.033 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Ammonium Laureth-2 Sulfate (UNII: 698O4Z48G6)	
Alcohol (UNII: 3K9958V90M)	
Isopropyl Alcohol (UNII: ND2M416302)	
Peg-25 Hydrogenated Castor Oil (UNII: 0ZNO9PJ9J)	
Polyquaternium-10 (125 Mpa.S At 2%) (UNII: L45WU8S981)	
Water (UNII: 059QF0K00R)	

Product Characteristics

Color	YELLOW (cloudy to transparent, cream to straw)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82392-100-04	1 in 1 CARTON		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M031	01/05/2021	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M031	01/05/2021	

Labeler - Oystershell Consumer Health Inc (117628199)

Revised: 12/2023

Oystershell Consumer Health Inc