

**PHENAZOPYRIDINE HYDROCHLORIDE 200 MG- phenazopyridine hydrochloride tablet**  
**GRAXCELL PHARMACEUTICAL, LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**Phenazopyridine Hydrochloride tablet, film coated - 200 mg, 70795-1252**

**INDICATIONS AND USAGE**

Phenazopyridine is indicated for the symptomatic relief of pain, burning, urgency, frequency, and other discomforts arising from irritation of the lower urinary tract mucosa caused by infection, trauma, surgery, endoscopic procedures, or the passage of sounds or catheters. The use of

Phenazopyridine HCl for relief of symptoms should not delay definitive diagnosis and treatment of causative conditions. Because it provides only symptomatic relief, prompt appropriate treatment of the cause of pain must be instituted and Phenazopyridine HCl should be discontinued when

symptoms are controlled. The analgesic action may reduce or eliminate the need for systemic analgesics or narcotics. It is, however, compatible with antibacterial therapy and can help to relieve pain and discomfort during the interval before antibacterial therapy controls the infection. Treatment of a urinary tract infection with Phenazopyridine HCl should not exceed two days because there is a lack of evidence that the combined administration of Phenazopyridine HCl and an antibacterial provides greater benefit than administration of the antibacterial alone after two days. (See DOSAGE AND ADMINISTRATION section.)

**DOSAGE AND ADMINISTRATION**

200 mg Tablets: Average adult dosage is one tablet 3 times a day after meals.

When used concomitantly with an antibacterial agent for the treatment of a urinary tract infection, the administration of Phenazopyridine HCl should not exceed 2 days.

**INACTIVE INGREDIENTS**

Phenazopyridine HCl Tablets, USP contains the following inactive ingredients: croscarmellose sodium, colloidal silicon dioxide, hydroxypropyl methyl cellulose, magnesium stearate, maize (corn starch) microcrystalline cellulose, polyethylene glycol, povidone and pregelatinized starch.



**GRAXCELL**  
PHARMACEUTICAL, LLC.

NDC 70795-1252-0

# Phenazopyridine Hydrochloride Oral Tablet, USP

Each tablet contains:  
Phenazopyridine HCL, USP.....200 mg

**200 mg**

**Rx Only  
100 Tablets**



Manufactured by:  
GRAXCELL PHARMACEUTICAL LLC  
136 Oak Drive Syosset, N.Y. 11791

Distributed by:  
GRAXCELL PHARMACEUTICAL LLC  
130 Knickerbocker Avenue Suite J,  
Bohemia, NY 11716  
P070 (REV:12 - 20)

## Phenazopyridine tablet, film coat

Graxcell Pharmaceuticals LLC


**Disclaimer:** This drug has not been found to be safe and effective, and this labeling has not been approved by FDA.

*For further information about unapproved drugs, visit the FDA website at: <https://www.fda.gov/drugs/development-research-and-manufacturing/unapproved-drugs>*

## (Phenazopyridine Hydrochloride Tablets, USP) Rx Only

**CAUTION:** Federal law prohibits dispensing without prescription.

**DESCRIPTION** Phenazopyridine Hydrochloride is a dark red to dark violet, odorless, slightly crystalline powder. It has a specific local anesthetic effect in the urinary tract, promptly relieving urinary discomfort and pain. It has the following structural formula:



C<sub>11</sub> H<sub>11</sub> N<sub>5</sub> • HCl

IV













# PHENAZOPYRIDINE HYDROCHLORIDE 200 MG

phenazopyridine hydrochloride tablet

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70795-1252
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENAZOPYRIDINE HYDROCHLORIDE (UNII: 0EWG668W17) (PHENAZOPYRIDINE - UNII:K2J09EMJ52)	PHENAZOPYRIDINE HYDROCHLORIDE	200 mg

## Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G17
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70795-1252-0	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/12/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/12/2020	

**Labeler** - GRAXCELL PHARMACEUTICAL, LLC (056556923)

Revised: 12/2021

GRAXCELL PHARMACEUTICAL, LLC