

ACETAMINOPHEN AND DIPHENHYDRAMINE HYDROCHLORIDE- acetaminophen and diphenhydramine hydrochloride tablet
Bi-Mart

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever
Diphenhydramine HCl 25 mg	Nighttime sleep aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- **adults and children 12 years and over:**take 2 caplets at bedtime; do not take more than 2 caplets of this product in 24 hours
- **children under 12 years:**do not use

Other information

- store between 20-25°C (68-77°F)
- DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Inactive ingredients

brilliant blue lake, colloidal silicon dioxide, indigo carmine lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions?

call toll free **1-844-912-4012**

PRINCIPAL DISPLAY PANEL-100 Count

NDC 37835-537-01

Compare to the active ingredients in Extra Strength Tylenol® PM*

EXTRA STRENGTH PAIN RELIEVER PM

Acetaminophen, Diphenhydramine HCl

Pain Reliever / Nighttime Sleep Aid

100 Caplets

*This product is not manufactured or distributed by the owner of the registered trademark Extra Strength Tylenol® PM.

Distributed by:

BI-MART

Eugene, OR 97402

Compare to the active ingredients in Extra Strength Tylenol® PM*

BI-MART NDC 37835-537-01

EXTRA STRENGTH PAIN RELIEVER PM

Non-Habit Forming

ACETAMINOPHEN
DIPHENHYDRAMINE HCl
Pain Reliever / Nighttime Sleep Aid

100 CAPLETS

0 71357 00674 8

Distributed by: BI-MART, Eugene, OR 97402

BI-MART UNCONVENTIONAL. MONITOR BACK LABELS FOR CHANGES. If you are not completely satisfied with your BI-MART purchase, return the unused portion and your entire purchase price will be cheerfully refunded.

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredients (in each caplet) Purpose
Acetaminophen 500 mg Pain reliever
Diphenhydramine HCl 25 mg Nighttime sleep aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on skin ■ in children under 12 years of age ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ glaucoma

Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin ■ taking

PEEL FOR DIRECTIONS 537-01-BM 02/25

LIFT HERE

Drug Facts (continued)

sedatives or tranquilizers
■ drowsiness will occur
■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)
■ adults and children 12 years and over: take 2 caplets at bedtime; do not take more than 2 caplets of this product in 24 hours
■ children under 12 years: do not use

Other information
■ Store between 20-25°C (68-77°F)

Inactive ingredients
Brilliant blue lake, colloidal silicon dioxide, indigo carmine lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Extra Strength Tylenol® PM

PRINCIPAL DISPLAY PANEL-250 Count

NDC 37835-537-25

Compare to the active ingredients in Extra Strength Tylenol® PM*

EXTRA STRENGTH PAIN RELIEVER PM

Acetaminophen, Diphenhydramine HCl

Pain Reliever / Nighttime Sleep Aid

250 Caplets

*This product is not manufactured or distributed by the owner of the registered trademark Extra Strength Tylenol® PM.

Distributed by:

BI-MART

Eugene, OR 97402

0 71357 00821 6

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
If you are not completely satisfied, you may return this product to the BI-MART store for a full refund. No receipt necessary. Price will be cheerfully refunded.

Compare to the active ingredients in Extra Strength Tylenol® PM*

VP VALUE PACK

BI-MART NDC 37835-537-25

EXTRA STRENGTH PAIN RELIEVER PM

Non-Habit Forming

**ACETAMINOPHEN
DIPHENHYDRAMINE HCl**

Pain Reliever / Nighttime Sleep Aid

250 CAPLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts	<p>Active ingredients (in each caplet)</p> <p>Acetaminophen 500 mg.....Pain reliever</p> <p>Diphenhydramine HCl 25 mg.....Nighttime sleep aid</p> <p>Uses: temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness</p> <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on skin</p> <ul style="list-style-type: none"> ■ in children under 12 years of age ■ if you have ever had an allergic reaction to this product or any of its ingredients <p>Ask a doctor before use if you have ■ liver disease ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland ■ glaucoma</p>
-------------------	---

▶ **PEEL FOR DIRECTIONS** 537-25-BM 02/25

LIFT HERE

Drug Facts (continued)	<p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers <p>When using this product ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ sleeplessness persists continuously for more than 2 weeks. ■ Insomnia may be a symptom of serious underlying medical illness. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed (see overdose warning) ■ adults and children 12 years and over: take 2 caplets at bedtime; do not take more than 2 caplets of this product in 24 hours ■ children under 12 years: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) <p>Inactive ingredients</p> <p>Brilliant blue lake, colloidal silicon dioxide, indigo carmine lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.</p> <p>Questions? call toll-free 1-844-912-4012</p> <p><small>*This product is not manufactured or distributed by the owner of the registered trademark Extra Strength Tylenol® PM</small></p>
-------------------------------	---

ACETAMINOPHEN AND DIPHENHYDRAMINE HYDROCHLORIDE

acetaminophen and diphenhydramine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-537
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE (UNII: 8GTS82S83M) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE	25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	APM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-537-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	
2	NDC:37835-537-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC Monograph Drug	M013	02/01/2025	
--------------------	------	------------	--

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
ELYSIUM PHARMACEUTICALS LIMITED		915664486	manufacture(37835-537)

Revised: 7/2025

Bi-Mart