

**MUCINEX FAST-MAX COLD, FLU AND SORE THROAT MAXIMUM STRENGTH-  
acetaminophen, dextromethorphan hydrobromide, guaifenesin, and  
phenylephrine hydrochloride capsule, liquid filled  
RB Health (US) LLC**

-----  
**Mucinex® Cold & Flu**

***Drug Facts***

<b><i>Active ingredients (in each liquid gel)</i></b>	<b><i>Purposes</i></b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

temporarily relieves these common cold and flu symptoms:

- minor aches and pains
- headache ■ sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not use more than directed**

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

## **Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat

## **Inactive ingredients**

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

## **Questions?**

### **1-866-MUCINEX (1-866-682-4639)**

You may also report side effects to this phone number.

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224

## **PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton**

Fast Dissolving Liquid Gels!

NDC 72854-127-16

MAXIMUM STRENGTH

Mucinex®

FAST-MAX®

COLD, FLU

& SORE THROAT

Acetaminophen – Pain Reliever/Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Guaifenesin – Expectorant

Phenylephrine HCl – Nasal Decongestant

✓ Controls Cough, Thins & Loosens Mucus

✓ Relieves Nasal & Chest Congestion

✓ Relieves Headache

✓ Reduces Fever

LIQUID GELS

(Liquid Filled Capsules)

Actual Size

DAY TIME

FOR AGES 12+

**MAXIMUM STRENGTH**

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

**Mucinex**  
**FAST-MAX**

COLD, FLU & SORE THROAT

**16 LIQUID GELS**

*Fast Dissolving Liquid Gels!*

**MAXIMUM STRENGTH** NDC 72854-127-16

**Mucinex**  
**FAST-MAX**

COLD, FLU & SORE THROAT

Dist. by RB Health (US)  
Paramaribo, NJ 07654-0224  
©2021 RB Health  
030121 3158654

For Cold & Flu Multi-Symptom relief for both Day and Night, try our **Maximum Strength Mucinex® FAST-MAX® Day/Night** liquid gel packs.

**MAXIMUM STRENGTH**  
COLD, FLU & SORE THROAT

Maximum Strength per 4-hour dose  
Do not take more than a total of  
12 liquid gels in a 24-hour period.  
Take only as directed.  
Keep carton for full information.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Acetaminophen – Pain Reliever/Fever Reducer**  
**Dextromethorphan HBr – Cough Suppressant**  
**Guaifenesin – Expectorant**  
**Phenylephrine HCl – Nasal Decongestant**

- ✓ Controls Cough, Thins & Loosens Mucus
- ✓ Relieves Nasal & Chest Congestion
- ✓ Relieves Headache
- ✓ Reduces Fever

Actual Size

**16 LIQUID GELS**  
**(Liquid Filled Capsules)**

DAY TIME  
FOR AGES 12+

LOT:

HEALTH • HYGIENE • HOME  
[www.mucinex.com](http://www.mucinex.com)  
Patents: [www.rb.com/patents](http://www.rb.com/patents)

3 6382458616 8

EXP.:

3158954



### Drug Facts

#### Active ingredients (in each liquid gel)

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

#### Purposes

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - headache
  - cough
  - minor aches and pains
  - sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

#### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

### Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

#### When using this product do not use more than directed

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

#### Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

#### Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

#### Questions? 1-866-MUCHEX (1-866-682-4639)

You may also report side effects to this phone number.

# MUCINEX FAST-MAX COLD, FLU AND SORE THROAT MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72854-127
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	24mm
<b>Flavor</b>		<b>Imprint Code</b>	AR01
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-127-	3 in 1 CARTON	07/26/2021	

1	16	2 III 1 CARTON	07/20/2021	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012		07/26/2021	

**Labeler -** RB Health (US) LLC (081049410)

Revised: 9/2022

RB Health (US) LLC