ROBAFEN DM- dextromethorphan hbr, guaifenes in solution Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major Pharmaceuticals Robafen® DM Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg

Purposes

Cough suppressant Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children	10 mL	
12 years and over	every 4 hours	
children under 12 years	do not use	

Other information

- Phenylketonurics:Contains Phenylalanine 30 mg per 10 mL
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

aspartame, benzoic acid, flavor, glycerin, hydroxyethyl cellulose, menthol, polyethylene glycol, purified water

Questions or comments?

1-800-616-2471

Principal Display Panel

ROBAFEN® DM COUGH

SUGAR-FREE

CLEAR

COUGH SUPPRESSANT (Dextromethorphan HBr)

EXPECTORANT (Guaifenesin)

Relieves:

Cough

Chest Congestion

Non-Drowsy


Specially Formulated For DIABETICS

COMPARE TO the active ingredients of ROBITUSSIN $^{\circledR}$ SUGAR-FREE COUGH + CONGESTION DM

FOR ADULTS

Ages 12 and Over

4 FL. OZ. (118 mL)



NDC 0904-6759-20

MAJOR°

NDC 0904-6759-20

MAJOR°

SUGAR-FREE

COUGH SUGAR-FREE

CLEAR

COUGH SUPPRESSANT (Dextromethorphan HBr) **EXPECTORANT** (Guaifenesin)

Relieves:

- □ Cough
- Chest Congestion

Non-Drowsy



COMPARE TO the active ingredients of ROBITUSSIN® SUGAR-FREE COUGH + CHEST CONGESTION DM*

FOR ADULTS Ages 12 and Over

4 FL. OZ. (118 mL)

CLEAR

COUGH SUPPRESSANT (Dextromethorphan HBr) **EXPECTORANT** (Guaifenesin)

Relieves:

- □ Cough
- Chest Congestion

Non-Drowsy

Specially Formulated For DIABETICS

COMPARE TO the active ingredients of ROBITUSSIN® SUGAR-FREE COUGH + CHEST CONGESTION DM*

FOR ADULTS Ages 12 and Over

4 FL. OZ. (118 mL)

Drug Facts

∄R∓RA∂US CONCH

Active ingredients (in each 10 mL) Dextromethorphan HBr, USP 20 mg.....

..Cough suppressant in, USP 200 mg...

MAJOR®

Purposes

GLEAR

- Uses temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tube

Warnings

Do not use if you are now taking a Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. this product.

Ask a doctor before use if you have cough that occurs with too much

phlegm (mucus)
cough that lasts or is chronic such as occurs with smoking, asthma, chronic

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

- Directions
 do not take more than 6 doses in any 24-hour period
 measure only with dosing cup provided
- keep dosing cup with product
 mL = milliliter
- this adult product is not intended for use in children under 12 years of age

Drug Facts (continued)

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- Phenylketonurics: Contains
 Phenylalanine 30 mg per 10 mL
 store at 20-25°C (68-77°F).
 Do not refrigerate.

Inactive ingredients aspartame, benzoic acid, flavor, glycerin, hydroxyethyl cellulose, menthol, polyethylene glycol, purified water

Questions or comments? 1-800-616-2471

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Sugar-Free Cough + Chest Congestion DM.


DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Distributed by MAJOR® PHARMACEUTICALS 17177 N Laurel Park Drive, Suite 233 Livonia, MI 48152 M-05 REV. 07/18 Re-Order No. 700959

Dosage Cup Included



www.StopMedicineAbuse.org





LOT NO.

EXP.

: 57826 SC C7

ROBAFEN DM

dextromethorphan hbr, guaifenesin solution

Product Information

Route of Administration

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0904-6759

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients		
Ingredient Name	Strength	
ASPARTAME (UNII: Z0H242BBR1)		
BENZOIC ACID (UNII: 8 SKN0 B0 MIM)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)		
WATER (UNII: 059QF0KO0R)		
MENTHOL (UNII: L7T10EIP3A)		

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

]	Packaging				
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0904-6759-20	1 in 1 CARTON	11/29/2018		
1	1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/29/2018	

Labeler - Major Pharmaceuticals (191427277)

Revised: 11/2018 Major Pharmaceuticals