

CIMETIDINE 200 ACID REDUCER- cimetidine tablet, film coated
Walgreen Company

Walgreen Co. Cimetidine 200 Drug Facts

Active ingredient (in each tablet)

Cimetidine 200 mg

Purpose

Acid reducer

Uses

- **relieves** heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking

- a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30minutes before** eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredient in Tagamet® HB 200

Cimetidine 200

CIMETIDINE TABLETS USP, 200 mg / ACID REDUCER

- Just one tablet relieves & prevents heartburn & acid indigestion
- Take before, during or after a meal

60 TABLETS (60 DOSES)

ACTUAL SIZE

Walgreens




Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Cimetidine 200

CIMETIDINE TABLETS USP, 200 mg / ACID REDUCER



Walgreens



Compare to the active ingredient in Tagamet® HB 200[†]

Cimetidine 200

CIMETIDINE TABLETS USP, 200 mg / ACID REDUCER

- Just one tablet relieves & prevents heartburn & acid indigestion
- Take before, during or after a meal

60

TABLETS
(60 DOSES)



ACTUAL SIZE

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING



ACTUAL SIZE

Walgreens

Cimetidine 200

CIMETIDINE TABLETS USP, 200 mg / ACID REDUCER

- Just one tablet relieves & prevents heartburn & acid indigestion
- Take before, during or after a meal

60 TABLETS (60 DOSES)

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

As the ingredient in each tablet

Purpose

Cimetidine 200 mg Acid reducer

NDC 0363-0022-72

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2022 Walgreen Co.

ITEM 491896 W10106-1124-L



3 11917 05411 7



PAPER BOX PLASTIC BOTTLE

how2recycle.info

W3ORG0223-F
REV-0824

02272 94 C7

Drug Facts

Active ingredient (in each tablet)

Cimetidine 200 mg.....Acid reducer

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- kidney disease
- liver disease

Ask a doctor or pharmacist before use if you are taking

- a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- you need to take this product for more than 14 days
- stomach pain continues
- your heartburn continues or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water **right before or any time up to 30 minutes before** eating food or drinking beverages that cause heartburn
 - do not take more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments?

1-800-719-9260

[†]Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

^{††}This product is not manufactured or distributed by Medtech Products Inc., distributor of Tagamet[®] HB 200.

CIMETIDINE 200 ACID REDUCER

cimetidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0022
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIMETIDINE (UNII: 80061L1WGD) (CIMETIDINE - UNII:80061L1WGD)	CIMETIDINE	200 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	L022
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0022-72	1 in 1 CARTON	04/14/2010	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075285	03/27/2001	

Labeler - Walgreen Company (008965063)

