

BLISTEX ENHANCEMENT SERIES- dimethicone, octinoxate, oxybenzone, avobenzone, and octocrylene

Blis tex Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Blis tex®

Enhancement Series®

DEEP RENEWAL®

Drug Facts

| <i>Active ingredients</i> | <i>Purpose</i> |
|----------------------------------|-----------------------|
| Avobenzone 3.0% (w/w) | Sunscreen |
| Dimethicone 2.0% (w/w) | Lip protectant |
| Octinoxate 7.5% (w/w) | Sunscreen |
| Octocrylene 2.7% (w/w) | Sunscreen |
| Oxybenzone 2.5% (w/w) | Sunscreen |

Uses

- temporarily protects and helps relieve chapped or cracked lips
- helps prevent sunburn

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure
- reapply at least every 2 hours
- use a water resistant sunscreen if swimming or sweating
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m.-2 p.m.
 - wear long sleeve shirts, pants, hats, and sunglasses
 - children under 6 months of age: Ask a doctor

Other information

- protect the product from excessive heat and direct sun

Inactive ingredients

atelocollagen, beeswax, butylene glycol, dipalmitoyl hydroxyproline, ethylhexyl palmitate, flavor, jojoba esters, lanolin alcohol, microcrystalline wax, octyldodecanol, pentaerythrityl tetraisostearate, petrolatum, phenoxyethanol, polybutene, polyethylene, PTFE, ricinus communis (castor) seed oil, saccharin, silica dimethyl silylate, sodium chondroitin sulfate, sodium hyaluronate, sorbic acid, squalane, tetrahexyldecyl ascorbate, tocopheryl acetate, ubiquinone

LIP VIBRANCE®

Drug Facts

| <i>Active ingredients</i> | <i>Purpose</i> |
|---------------------------|----------------|
| Dimethicone 2.0% (w/w) | Lip protectant |
| Octinoxate 7.5% (w/w) | Sunscreen |
| Oxybenzone 2.5% (w/w) | Sunscreen |

Uses

- temporarily protects and helps relieve chapped or cracked lips
- helps prevent sunburn

Warnings

Skin Cancer/Skin Aging Alert

Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure
- reapply at least every 2 hours
- use a water resistant sunscreen if swimming or sweating
- children under 6 months of age: Ask a doctor

Other information

- protect the product from excessive heat and direct sun

Inactive ingredients

arachidyl alcohol, arachidyl glucoside, behenyl alcohol, bis-diglyceryl polyacyladipate-2, butylene glycol, butyrospermum parkii (shea butter), D&C red no. 6, D&C red no. 7, ethylhexyl palmitate, flavors, hydrogenated polyisobutene, iron oxides, isostearoyl hydrolyzed silk, mica, microcrystalline wax, mineral oil, ozokerite, panthenyl ethyl ether, paraffin, pentaerythrityl tetraethylhexanoate, petrolatum, phenoxyethanol, phenyl trimethicone, polybutene, polyethylene, PPG-5-ceteth-20, ricinus

communis (castor) seed oil, saccharin, silica dimethyl silylate, sodium hyaluronate, titanium dioxide, tocopheryl acetate, vitis vinifera (grape) seed oil

SILK & SHINE®

Drug Facts

| <i>Active ingredients</i> | <i>Purpose</i> |
|----------------------------------|-----------------------|
| Dimethicone 2.0% (w/w) | Lip protectant |
| Octinoxate 7.5% (w/w) | Sunscreen |
| Oxybenzone 2.5% (w/w) | Sunscreen |

Uses

- temporarily protects and helps relieve chapped or cracked lips
- helps prevent sunburn

Warnings

Skin Cancer/Skin Aging Alert

Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure
- reapply at least every 2 hours
- use a water resistant sunscreen if swimming or sweating
- children under 6 months of age: Ask a doctor

Other information

- protect the product from excessive heat and direct sun

Inactive ingredients

arachidyl alcohol, arachidyl glucoside, behenyl alcohol, bis-diglyceryl polyacyladipate-2, flavors, hydrogenated polyisobutene, isostearoyl hydrolyzed silk, mica, microcrystalline wax, mineral oil, ozokerite, panthenyl ethyl ether, paraffin, pentaerythrityl tetraethylhexanoate, petrolatum, phenoxyethanol, phenyl trimethicone, polybutene, PPG-5-Ceteth-20, saccharin, titanium dioxide, tocopheryl acetate

PRINCIPAL DISPLAY PANEL - 3 Stick Blister Pack

NDC 10157-9858-1

Blistex®
LIP PROTECTANT/SUNSCREEN

ENHANCEMENT SERIES®
youthful • beautiful • kissable

BUY 2 GET 1 FREE

BROAD SPECTRUM
SPF15

SPF15

SPF15

3 Sticks Net Wt. 0.13 oz. (3.69 g) Each



DEEP RENEWAL. The anti-aging lip treatment with coenzyme Q-10, clinically proven to visibly reduce fine lines and wrinkles in just four weeks.

| | | | | | | | | | | | | |
|--|-----------------------|-----------|------------------------|----------------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|---|--|
| <p>Drug Facts</p> <p>Active Ingredients</p> <table border="0"> <tr> <td>Avobenzone 3.0% (w/w)</td> <td>Sunscreen</td> </tr> <tr> <td>Dimethicone 2.0% (w/w)</td> <td>Lip protectant</td> </tr> <tr> <td>Octinoxate 7.5% (w/w)</td> <td>Sunscreen</td> </tr> <tr> <td>Octocrylene 2.7% (w/w)</td> <td>Sunscreen</td> </tr> <tr> <td>Oxybenzone 2.5% (w/w)</td> <td>Sunscreen</td> </tr> </table> <p>Purpose</p> <p>• temporarily protects and helps relieve chapped or cracked lips • helps prevent sunburn</p> <p>Warnings</p> <p>For external use only</p> <p>Do not use on damaged or broken skin</p> <p>When using this product keep out of eyes. Rinse with water to remove.</p> | Avobenzone 3.0% (w/w) | Sunscreen | Dimethicone 2.0% (w/w) | Lip protectant | Octinoxate 7.5% (w/w) | Sunscreen | Octocrylene 2.7% (w/w) | Sunscreen | Oxybenzone 2.5% (w/w) | Sunscreen | <p>Stop use and ask a doctor if rash occurs.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <p>• apply liberally 15 minutes before sun exposure • reapply at least every 2 hours • use a water resistant sunscreen if swimming or sweating • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m.-2 p.m. • wear long sleeve shirts, pants, hats, and sunglasses • children under 6 months of age: Ask a doctor</p> | <p>Other Information</p> <p>• protect the product from excessive heat and direct sun</p> <p>Inactive Ingredients</p> <p>atelo collagen, beeswax, butylene glycol, dipalmitoyl hydroxyproline, ethylhexyl palmitate, flavor, jojoba esters, lanolin alcohol, microcrystalline wax, octyldodecanol, pentaerythrityl tetraoctadecanoate, petrolatum, phenoxethanol, polybutene, polyethylene, PTE, nicotinic acid, (castor) seed oil, saccharin, silica dimethyl silylate, sodium chondroitin sulfate, sodium hyaluronate, sorbic acid, squalane, tetrahydroxybutyl ascorbate, tocopheryl acetate, ubiquinone</p> |
| Avobenzone 3.0% (w/w) | Sunscreen | | | | | | | | | | | |
| Dimethicone 2.0% (w/w) | Lip protectant | | | | | | | | | | | |
| Octinoxate 7.5% (w/w) | Sunscreen | | | | | | | | | | | |
| Octocrylene 2.7% (w/w) | Sunscreen | | | | | | | | | | | |
| Oxybenzone 2.5% (w/w) | Sunscreen | | | | | | | | | | | |

LIP VIBRANCE. Beautiful, buildable color and shimmer that really lasts, plus shea butter, grapeseed oil and vitamin E.

| | | | | | | | | |
|---|------------------------|----------------|-----------------------|-----------|-----------------------|-----------|---|--|
| <p>Drug Facts</p> <p>Active Ingredients</p> <table border="0"> <tr> <td>Dimethicone 2.0% (w/w)</td> <td>Lip protectant</td> </tr> <tr> <td>Octinoxate 7.5% (w/w)</td> <td>Sunscreen</td> </tr> <tr> <td>Oxybenzone 2.5% (w/w)</td> <td>Sunscreen</td> </tr> </table> <p>Purpose</p> <p>• temporarily protects and helps relieve chapped or cracked lips • helps prevent sunburn</p> <p>Warnings</p> <p>Skin Cancer/Skin Aging Alert: Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, not skin cancer or early skin aging. For external use only</p> | Dimethicone 2.0% (w/w) | Lip protectant | Octinoxate 7.5% (w/w) | Sunscreen | Oxybenzone 2.5% (w/w) | Sunscreen | <p>Do not use on damaged or broken skin</p> <p>When using this product keep out of eyes. Rinse with water to remove.</p> <p>Stop use and ask a doctor if rash occurs</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <p>• apply liberally 15 minutes before sun exposure • reapply at least every 2 hours • use a water resistant sunscreen if swimming or sweating • children under 6 months of age: Ask a doctor</p> <p>Other Information</p> <p>• protect the product from excessive heat and direct sun</p> | <p>Inactive Ingredients</p> <p>arachidyl alcohol, arachidyl glucoside, behenyl alcohol, bis-diglyceryl polyacrylate-2, butylene glycol, butyrospermum parkii (shea butter), D&C red no. 6, D&C red no. 7, ethylhexyl palmitate, flavors, hydrogenated polyisobutene, iron oxides, isostearyl hydrolyzed silk, mica, microcrystalline wax, mineral oil, ozokerite, paraffin, pentaerythrityl tetraethylhexanoate, petrolatum, phenoxethanol, phenyl trimethicone, polybutene, polyethylene, PPG-5-ceteth-20, nicotinic acid, (castor) seed oil, saccharin, silica dimethyl silylate, sodium hyaluronate, titanium dioxide, tocopheryl acetate, vitis vinifera (grape) seed oil</p> |
| Dimethicone 2.0% (w/w) | Lip protectant | | | | | | | |
| Octinoxate 7.5% (w/w) | Sunscreen | | | | | | | |
| Oxybenzone 2.5% (w/w) | Sunscreen | | | | | | | |

SILK & SHINE. Enhances lips' natural radiance, leaves a smoother lip feel, and keeps lips hydrated...with real silk extracts and vitamins.

| | | | | | | | | |
|---|------------------------|----------------|-----------------------|-----------|-----------------------|-----------|---|--|
| <p>Drug Facts</p> <p>Active Ingredients</p> <table border="0"> <tr> <td>Dimethicone 2.0% (w/w)</td> <td>Lip protectant</td> </tr> <tr> <td>Octinoxate 7.5% (w/w)</td> <td>Sunscreen</td> </tr> <tr> <td>Oxybenzone 2.5% (w/w)</td> <td>Sunscreen</td> </tr> </table> <p>Purpose</p> <p>• temporarily protects and helps relieve chapped or cracked lips • helps prevent sunburn</p> <p>Warnings</p> <p>Skin Cancer/Skin Aging Alert: Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, not skin cancer or early skin aging. For external use only</p> | Dimethicone 2.0% (w/w) | Lip protectant | Octinoxate 7.5% (w/w) | Sunscreen | Oxybenzone 2.5% (w/w) | Sunscreen | <p>Do not use on damaged or broken skin</p> <p>When using this product keep out of eyes. Rinse with water to remove.</p> <p>Stop use and ask a doctor if rash occurs</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <p>• apply liberally 15 minutes before sun exposure • reapply at least every 2 hours • use a water resistant sunscreen if swimming or sweating • children under 6 months of age: Ask a doctor</p> <p>Other Information</p> <p>• protect the product from excessive heat and direct sun</p> | <p>Inactive Ingredients</p> <p>arachidyl alcohol, arachidyl glucoside, behenyl alcohol, bis-diglyceryl polyacrylate-2, flavors, hydrogenated polyisobutene, isostearyl hydrolyzed silk, mica, microcrystalline wax, mineral oil, ozokerite, paraffin, pentaerythrityl tetraethylhexanoate, petrolatum, phenoxethanol, phenyl trimethicone, polybutene, PPG-5-Ceteth-20, saccharin, titanium dioxide, tocopheryl acetate</p> |
| Dimethicone 2.0% (w/w) | Lip protectant | | | | | | | |
| Octinoxate 7.5% (w/w) | Sunscreen | | | | | | | |
| Oxybenzone 2.5% (w/w) | Sunscreen | | | | | | | |

#41221 ©2014 Blistex Inc. P.O. Box 5382, Oak Brook, IL 60522-5382

SATISFACTION GUARANTEED

Blistex

Card is 100% Recyclable.

0 41388 00435 8

BLISTEX ENHANCEMENT SERIES

dimethicone, octinoxate, oxybenzone, avobenzone, and octocrylene kit

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:10 157-9858

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:10157-9858-1 | 1 in 1 BLISTER PACK | | |

Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 CYLINDER | 3.69 g |
| Part 2 | 1 CYLINDER | 3.69 g |
| Part 3 | 1 CYLINDER | 3.69 g |

Part 1 of 3

BLISTEX LIP VIBRANCE

dimethicone, octinoxate, and oxybenzone stick

Product Information

Route of Administration TOPICAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| Dimethicone (UNII: 92RU3N3Y1O) (Dimethicone - UNII:92RU3N3Y1O) | Dimethicone | 2 g in 100 g |
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 7.5 g in 100 g |
| Oxybenzone (UNII: 95OOS7VE0Y) (Oxybenzone - UNII:95OOS7VE0Y) | Oxybenzone | 2.5 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| arachidyl alcohol (UNII: 1QR1QRA9BU) | |
| ceresin (UNII: Q1LS2UJO3A) | |
| paraffin (UNII: I9O0E3H2ZE) | |
| microcrystalline wax (UNII: XOF597Q3KY) | |
| petrolatum (UNII: 4T6H12BN9U) | |
| mineral oil (UNII: T5L8T28FGP) | |
| arachidyl glucoside (UNII: 6JVV35JOOJ) | |
| docosanol (UNII: 9G1OE216XY) | |
| butylene glycol (UNII: 3XUS85K0RA) | |
| shea butter (UNII: K49155WL9Y) | |
| D&C red no. 7 (UNII: ECW0LZ41X8) | |
| D&C red no. 6 (UNII: 481744A14O) | |
| ethylhexyl palmitate (UNII: 2865993309) | |
| phenoxyethanol (UNII: HIE492ZZ3T) | |
| mica (UNII: V8A1AW0880) | |

| |
|--|
| panthenyl ethyl ether (UNII: F4WMF8NX3B) |
| pentaerythrityl tetraethylhexanoate (UNII: XJ7052W897) |
| high density polyethylene (UNII: UG00KM4WR7) |
| PPG-5-ceteth-20 (UNII: 4AAN25P8P4) |
| ferric oxide red (UNII: 1K09F3G675) |
| titanium dioxide (UNII: 15FIX9V2JP) |
| phenyl trimethicone (UNII: DR0K5NOJ4R) |
| grape seed oil (UNII: 930MLC8XGG) |
| castor oil (UNII: D5340Y2I9G) |
| sandalwood (UNII: 3641YW25N2) |
| saccharin (UNII: FST467XS7D) |
| .alpha.-tocopherol acetate (UNII: 9E8X80D2L0) |
| silica dimethyl silylate (UNII: EU2PSP0G0W) |
| hyaluronate sodium (UNII: YSE9PPT4TH) |
| Polybutene (1400 MW) (UNII: 1NA5AO9GH7) |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|----------------------|----------------------|--------------------|
| 1 | | 3.69 g in 1 CYLINDER | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part347 | 05/31/2013 | |

Part 2 of 3

BLISTEX SILK AND SHINE

dimethicone, octinoxate, and oxybenzone stick

Product Information

| | |
|-------------------------|---------|
| Route of Administration | TOPICAL |
|-------------------------|---------|

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| Dimethicone (UNII: 92RU3N3Y1O) (Dimethicone - UNII:92RU3N3Y1O) | Dimethicone | 2 g in 100 g |
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 7.5 g in 100 g |
| Oxybenzone (UNII: 95OOS7VE0Y) (Oxybenzone - UNII:95OOS7VE0Y) | Oxybenzone | 2.5 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------|----------|
| arachidyl alcohol (UNII: 1QR1QRA9BU) | |

| |
|--|
| arachidyl glucoside (UNII: 6JVV35J00J) |
| docosanol (UNII: 9G1OE216XY) |
| mica (UNII: V8A1AW0880) |
| microcrystalline wax (UNII: XOF597Q3KY) |
| mineral oil (UNII: T5L8T28FGP) |
| panthenyl ethyl ether (UNII: F4WMF8NX3B) |
| paraffin (UNII: I9O0E3H2ZE) |
| pentaerythrityl tetraethylhexanoate (UNII: XJ7052W897) |
| petrolatum (UNII: 4T6H12BN9U) |
| phenoxyethanol (UNII: HIE492ZZ3T) |
| phenyl trimethicone (UNII: DR0K5NOJ4R) |
| PPG-5-Ceteth-20 (UNII: 4AAN25P8P4) |
| saccharin (UNII: FST467XS7D) |
| titanium dioxide (UNII: 15FIX9V2JP) |
| sandalwood (UNII: 3641YW25N2) |
| .alpha.-tocopherol acetate (UNII: 9E8X80D2L0) |
| Ceresin (UNII: Q1LS2UJO3A) |
| Polybutene (1400 MW) (UNII: 1NA5AO9GH7) |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|----------------------|----------------------|--------------------|
| 1 | | 3.69 g in 1 CYLINDER | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part347 | 11/01/2012 | |

Part 3 of 3

BLISTEX DEEP RENEWAL

avobenzene, octinoxate, octocrylene, oxybenzone, and dimethicone stick

Product Information

| | |
|-------------------------|---------|
| Route of Administration | TOPICAL |
|-------------------------|---------|

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| Dimethicone (UNII: 92RU3N3Y1O) (Dimethicone - UNII:92RU3N3Y1O) | Dimethicone | 2 g in 100 g |
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 7.5 g in 100 g |
| Oxybenzone (UNII: 95OOS7VE0Y) (Oxybenzone - UNII:95OOS7VE0Y) | Oxybenzone | 2.5 g in 100 g |
| Avobenzene (UNII: G63QQF2NOX) (Avobenzene - UNII:G63QQF2NOX) | Avobenzene | 3 g in 100 g |
| Octocrylene (UNII: 5A68WGF6WM) (Octocrylene - UNII:5A68WGF6WM) | Octocrylene | 2.7 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| yellow wax (UNII: 2ZA36H0S2V) | |
| butylene glycol (UNII: 3XUS85K0RA) | |
| dipalmitoyl hydroxyproline (UNII: E6AHA53N1H) | |
| ethylhexyl palmitate (UNII: 2865993309) | |
| lanolin alcohols (UNII: 884C3FA9HE) | |
| microcrystalline wax (UNII: XOF597Q3KY) | |
| octyldodecanol (UNII: 461N1O614Y) | |
| pentaerythrityl tetraistearate (UNII: 9D7IK5483F) | |
| petrolatum (UNII: 4T6H12BN9U) | |
| phenoxyethanol (UNII: HIE492ZZ3T) | |
| high density polyethylene (UNII: UG00KM4WR7) | |
| polytetrafluoroethylene (UNII: E1NC1JVS3O) | |
| castor oil (UNII: D5340Y2I9G) | |
| saccharin (UNII: FST467XS7D) | |
| silica dimethyl silylate (UNII: EU2PSP0G0W) | |
| hyaluronate sodium (UNII: YSE9PPT4TH) | |
| sorbic acid (UNII: X045WJ989B) | |
| squalane (UNII: GW89575KF9) | |
| tetrahexyldecyl ascorbate (UNII: 9LBV3F07AZ) | |
| Polybutene (1400 MW) (UNII: 1NA5AO9GH7) | |
| MARINE COLLAGEN, SOLUBLE (UNII: 8JC99XGU4W) | |
| CHONDROITIN SULFATE SODIUM (SHARK) (UNII: Q75WVO004L) | |
| UBIDECARENONE (UNII: EJ27X76M46) | |
| SIMMONDSIA CHINENSIS SEED WAX (UNII: 47X6Y9FJJ2) | |
| .alpha.-tocopherol acetate (UNII: 9E8X80D2L0) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|----------------------|----------------------|--------------------|
| 1 | | 3.69 g in 1 CYLINDER | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part347 | 11/01/2012 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part347 | 05/15/2014 | |

Labeler - Blistex Inc. (005126354)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------|---------|-----------|--------------------------|
| Blistex Inc. | | 005126354 | MANUFACTURE(10 157-9858) |

Revised: 4/2014

Blistex Inc.