

SOMXL- salicylic acid solution
WR Group Skin Care Ltd

Drug Facts

Active ingredient

Salicylic acid 17% (w/w)

Purpose

Wart remover

Use

For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance on the surface.

Warnings

For external use only

If product gets into eyes, flush with water for 15-minutes. Avoid inhaling vapors.

Do not use

- on irritated skin
- on any area that is infected or reddened
- if you are diabetic or if you have poor blood circulation
- piles
- birthmarks
- warts with hair growing from them
- on moles
- warts on the face or mucous membranes

Stop use and ask a doctor if

discomfort persists

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Wash affected area (May soak wart in warm water for 15-minutes if desired).
- Dry area thoroughly.
- Apply small amount with applicator to cover each wart. Let dry.
- Repeat this procedure once or twice daily as needed for up to 12-weeks.

Inactive ingredients

castor oil, ethyl lactate, flexible collodion, polybutene, (Contains alcohol 18% v/v and ether 53% v/v from flexible collodion.)

Package Labeling:

Net wt. 0.5 oz (15 g)

SOMXL

Drug Facts

Active ingredient **Purpose**
Salicylic acid 17% (w/w) Wart remover

Use For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance on the surface.

Warnings

For external use only
If product gets into eyes, flush with water for 15-minutes. Avoid inhaling vapors.

Do not use ■ on irritated skin
■ on any area that is infected or reddened
■ if you are diabetic or if you have poor blood circulation ■ piles ■ birthmarks
■ warts with hair growing from them ■ on moles
■ warts on the face or mucous membranes

Stop use and ask a doctor if discomfort persists

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

■ Wash affected area (May soak wart in warm water for 15-minutes if desired).
■ Dry area thoroughly.
■ Apply small amount with applicator to cover each wart. Let dry.
■ Repeat this procedure once or twice daily as needed for up to 12-weeks.

Inactive ingredients

castor oil, ethyl lactate, flexible collodion, polybutene. (Contains alcohol 18% v/v and ether 53% v/v from flexible collodion.)

DISTRIBUTED BY WR GROUP SKIN CARE LTD
PO BOX 13867, CHRISTCHURCH, NEW ZEALAND

Net wt. 0.5 oz (15 g)



Wart Remover

Net wt. 0.5 oz (15 g)



Wart Remover

Net wt. 0.5 oz (15 g)



Wart Remover

SOMXL[®]

Net wt. 0.5 oz (15 g)

SOMXL

salicylic acid solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70463-000
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	170 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL LACTATE (UNII: F3P750VW8I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70463-000-01	1 in 1 BOX	08/01/2024	
1		15 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	08/01/2024	

Labeler - WR Group Skin Care Ltd (592719822)

Revised: 8/2024

WR Group Skin Care Ltd