

**EXTRA STRENGTH TYLENOL COLD PLUS FLU MULTI-ACTION DAY/NIGHT-
acetaminophen, pseudoephedrine hydrochloride, dextromethorphan
hydrobromide, and chlorpheniramine maleate
Kenvue Brands LLC**

Extra Strength TYLENOL Cold + Flu Multi-Action

Day/Night

TYLENOL® COLD + FLU MULTI-ACTION DAY

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Pseudoephedrine HCl 30 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 6 hours while symptoms last▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20 - 25°C (68 - 77°F).
- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, hypromellose, magnesium stearate, microcrystalline cellulose, powdered cellulose, pregelatinized starch, propylene glycol, sodium starch glycolate, titanium dioxide

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

TYLENOL® COLD + FLU MULTI-ACTION NIGHT

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 15 mg	Cough suppressant
Pseudoephedrine HCl 30 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - cough

- nasal congestion
- sinus congestion and pressure
- sneezing and runny nose
- helps clear nasal passages
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 6 hours while symptoms last▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20 - 25°C (68 - 77°F).

- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, FD&C blue No.1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-374-01

Extra Strength TYLENOL®

FOR ADULTS

COLD + FLU MULTI-ACTION

Acetaminophen

Dextromethorphan HBr, Pseudoephedrine HCl

Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant

DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION

Actual Size

16 CAPLETS;

Acetaminophen

Chlorpheniramine Maleate, Dextromethorphan HBr, Pseudoephedrine HCl

Pain Reliever-Fever Reducer, Antihistamine, Cough Suppressant, Nasal Decongestant

NIGHT

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH • NASAL CONGESTION
- RUNNY NOSE

Actual Size

8 CAPLETS; TOTAL 24 CAPLETS





How can we help?
1-877-895-3665

30051444

NDC 50580-374-01

Extra Strength

TYLENOL[®]
FOR ADULTS

COLD + FLU MULTI-ACTION

Acetaminophen

Dextromethorphan HBr, Pseudoephedrine HCl
Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant

Acetaminophen

Chlorpheniramine Maleate, Dextromethorphan HBr, Pseudoephedrine HCl
Pain Reliever-Fever Reducer, Antihistamine, Cough Suppressant, Nasal Decongestant



- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION

DAY



Actual Size

16 CAPLETS;

NIGHT



Actual Size

8 CAPLETS; TOTAL 24 CAPLETS

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH + NASAL CONGESTION
- RUNNY NOSE

Extra Strength
TYLENOL[®]
COLD + FLU MULTI-ACTION

DO NOT USE IF BLISTER UNIT IS TORN OR BROKEN.

Made in Canada
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©18/10/2021

Drug Facts (continued)

■ pain, nasal congestion or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)
adults and children
■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
children under 12 years
■ ask a doctor
■ do not use for more than 10 days unless directed by a doctor

Other information
■ store between 20°-25°C (68°-77°F)
■ do not use if blister unit is torn or broken

Inactive ingredients carmelum, FD&C blue #1, aluminum lake, titanium dioxide, hydroxyethylcellulose, polyethylene glycol, powdered cellulose, pregelatinized starch, sodium starch glycolate, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose.

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)
adults and children
■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
children under 12 years
■ ask a doctor
■ do not use for more than 10 days unless directed by a doctor

Other information
■ store between 20°-25°C (68°-77°F)
■ do not use if blister unit is torn or broken

Inactive ingredients carmelum, magnesium stearate, microcrystalline cellulose, polydextrose, starch glycolate, titanium dioxide

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

TYLENOL® COLD + FLU MULTI-ACTION NIGHT

▲ **Stop use and ask a doctor if** ■ nervousness, dizziness, or sleepiness occur
■ be careful when driving a motor vehicle or operating machinery
■ avoid alcoholic drinks
■ alcohol, sedatives and tranquilizers may increase drowsiness
■ excitability may occur, especially in children ■ marked drowsiness may occur

When using this product ■ do not exceed recommended dose
■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
■ cough that occurs with too much phlegm (mucus)
■ persistent or chronic cough such as occurs with smoking, asthma or emphysema
■ trouble urinating due to an enlarged prostate gland
■ diabetes ■ heart disease ■ high blood pressure ■ thyroid disease

Ask a doctor before use if you have
■ if you have ever had an allergic reaction to this product or any of its ingredients

TYLENOL® COLD + FLU MULTI-ACTION DAY

▲ **Stop use and ask a doctor if** ■ nervousness, dizziness, or sleepiness occur
■ pain, nasal congestion or cough gets worse or lasts more than 7 days


When using this product do not exceed recommended dose
■ taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
■ persistent or chronic cough such as occurs with smoking, asthma or emphysema
■ trouble urinating due to an enlarged prostate gland
■ diabetes ■ heart disease ■ high blood pressure

Ask a doctor before use if you have
■ if you have ever had an allergic reaction to this product or any of its ingredients

0045014924 4
3

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



Important: Read all product information before using. Keep this box for important information. Do not take the day and night caplets at the same time. Do not take more than a total of 6 caplets in a 24-hour period. Take only as directed.

TYLENOL® COLD + FLU MULTI-ACTION DAY

TYLENOL® COLD + FLU MULTI-ACTION NIGHT

Drug Facts
Active ingredients (in each caplet)
 Acetaminophen 500 mg
 Pain reliever/fever reducer
 Dextromethorphan HBr 15 mg
 Cough suppressant
 Pseudoephedrine HCl 30 mg
 Nasal decongestant

Uses
 Temporarily relieves these common cold/flu symptoms:
 ■ minor aches and pains ■ sore throat ■ headach
 ■ nasal congestion ■ sinus congestion and pressure
 ■ temporarily reduces fever
 ■ helps clear nasal passages
 ■ provides nasal and sinus drainage

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin redness ■ blisters ■ rash
 ■ 3 or more alcoholic drinks every day while using this product

Drug Facts
Active ingredients (in each caplet)
 Acetaminophen 500 mg
 Pain reliever/fever reducer
 Chlorpheniramine maleate 2 mg
 Antihistamine
 Dextromethorphan HBr 15 mg
 Cough suppressant
 Pseudoephedrine HCl 30 mg
 Nasal decongestant

Uses
 Temporarily relieves these common cold/flu symptoms:
 ■ minor aches and pains ■ sore throat ■ headach
 ■ nasal congestion ■ sinus congestion and pressure
 ■ temporarily reduces fever
 ■ helps clear nasal passages

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin redness ■ blisters ■ rash
 ■ 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, psychiatric or emotional conditions, or Parkinson's disease), or if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, psychiatric or emotional conditions, or Parkinson's disease), or drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, psychiatric or emotional conditions, or Parkinson's disease), or if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, psychiatric or emotional conditions, or Parkinson's disease), or if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist before taking this product.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, psychiatric or emotional conditions, or Parkinson's disease), or if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist before taking this product.

EXTRA STRENGTH TYLENOL COLD PLUS FLU MULTI-ACTION DAY/NIGHT

acetaminophen, pseudoephedrine hydrochloride, dextromethorphan hydrobromide, and chlorpheniramine maleate kit

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-374
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-374-01	1 in 1 PACKAGE	06/21/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	2 BLISTER PACK	8

Part 1 of 2

EXTRA STRENGTH TYLENOL COLD PLUS FLU MULTI-ACTION DAY

acetaminophen, pseudoephedrine hydrochloride, and dextromethorphan hydrobromide tablet, film coated

Product Information

Item Code (Source)	NDC:50580-344
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	TY;COLD;1408
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/21/2021	

Part 2 of 2

EXTRA STRENGTH TYLENOL COLD PLUS FLU MULTI-ACTION NIGHT

acetaminophen, chlorpheniramine maleate, pseudoephedrine hydrochloride, and dextromethorphan hydrobromide tablet, film coated

Product Information

Item Code (Source)	NDC:50580-286
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	TY;COLD;1407
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/21/2021	

Labeler - Kenvue Brands LLC (118772437)