

**NASAL- oxymetazoline hydrochloride spray, metered**  
**Rite Aid Corporation**

-----  
**Rite Aid Corporation Nasal Spray Drug Facts**

**Active ingredient**

Oxymetazoline hydrochloride 0.05%

**Purpose**

Nasal decongestant

**Uses**

- temporarily relieves nasal congestion due to:
- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

**Warnings**

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

- **do not use more than directed**
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

**Stop use and ask a doctor if**

symptoms persist

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

**To Use:** Shake well before use. Hold white tabs, SQUEEZE grooved area of cap FIRMLY and turn counter clockwise. Before using the first time, prime metered pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

**Other information**

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

**Inactive ingredients**

benzalkonium chloride solution, benzyl alcohol, camphor, dibasic sodium phosphate, edetate disodium, eucalyptol, menthol, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

PLUS MENTHOL

Compare to the active ingredient of Afrin<sup>®</sup> No Drip Severe Congestion

SEVERE CONGESTION

NASAL SPRAY

NASAL DECONGESTANT

OXYMETAZOLINE HCl

NO DRIP

Fast, powerful congestion relief

For colds & allergies

12 HOUR

1 FL OZ (30 mL)

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

DISTRIBUTED BY:  
RITE AID  
200 NEUBERRY COMMONS  
EITERS, PA 17319  
www.riteaid.com

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

**PLUS MENTHOL** NDC 11822-6378-1

Compare to the active ingredient of Afrin® No Drip Severe Congestion\*

**SEVERE CONGESTION**

**NASAL SPRAY**

**NASAL DECONGESTANT**

**OXYMETAZOLINE HCl**

**NO DRIP**

Fast, powerful congestion relief  
For colds & allergies

**12 HOUR**

**1 FL OZ (30 mL)**

**COOL AREA**

7J810 83 C2

**Drug Facts**

**Active ingredient** Purpose  
Oxymetazoline hydrochloride 0.05%...Nasal decongestant

**Uses**

- temporarily relieves nasal congestion due to:
  - common cold
  - hay fever
  - upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

**Warnings**

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

- do not use more than directed
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

**To Use:** Shake well before use. Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Before using the first time, prime metered pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

**Other information**

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

**Drug Facts (continued)**

**Inactive ingredients**  
benzalkonium chloride solution, benzyl alcohol, camphor, dibasic sodium phosphate, edetate disodium, eucalyptol, menthol, microcrystalline cellulose and carboxymethylcellulose sodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

**Questions or comments?**  
1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Afrin® No Drip Severe Congestion.

**How to use:**  
Hold white tabs, **SQUEEZE** grooved area of cap **FIRMLY** and turn counter clockwise. Secure cap after use.

**SQUEEZE** grooves **FIRMLY** while turning

Hold white tabs **FIRMLY**

**SEVERE CONGESTION** NDC 11822-6378-1  
**NASAL SPRAY**  
**NASAL DECONGESTANT**  
OXYMETAZOLINE HCl  
**NO DRIP**  
For colds & allergies  
**12 HOUR**  
1 FL OZ (30 mL)  
**ACTUAL SIZE**

# NASAL

oxymetazoline hydrochloride spray, metered

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-6378
<b>Route of Administration</b>	NASAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	.05 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM</b> (UNII: GR686LBA74)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>EUCALYPTOL</b> (UNII: RV6J6604TK)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2S W)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	

## Product Characteristics

<b>Color</b>	WHITE (off white)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-6378-1	1 in 1 CARTON	12/05/2018	
1		30 mL in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	12/05/2018	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 8/2024

Rite Aid Corporation