

**GOOD NEIGHBOR PHARMACY CHILDRENS IBUPROFEN- ibuprofen suspension**  
**Amerisource Bergen**

-----  
**Amerisource Bergen Children's Ibuprofen Oral Suspension Drug Facts**

**Active ingredient (in each 5 mL)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever,

headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- give with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

## Dosing Chart

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)**</b>
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\*or as directed by a doctor

## Other information

- each 5 mL contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze

## Inactive ingredients

anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

## Questions or comments?

1-800-719-9260

## Principal Display Panel

FOR AGES 2 TO 11 YEARS

children's

Ibuprofen

Oral Suspension

100 mg per 5 mL

pain reliever/fever reducer (NSAID)

Lasts up to 8 hours

Grape Flavor

Alcohol Free

See New Warnings

4 fl oz (120 mL)



NDC 48122-110-26



FOR AGES  
2 TO 11 YEARS

children's  
**Ibuprofen**

Oral Suspension

100 mg per 5 mL

pain reliever/  
fever reducer (NSAID)

Lasts up to  
8 hours

Grape Flavor

Alcohol Free

See New  
Warnings

4 fl oz  
(120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

**Drug Facts**

**Active ingredient** *Purposes*  
(in each 5 mL)  
Ibuprofen 100 mg (NSAID)\*.....Pain reliever/  
fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily:  
relieves minor aches and pains due to the  
common cold, flu, sore throat, headache  
and toothache  
reduces fever


**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe  
allergic reaction, especially in people allergic  
to aspirin. Symptoms may include:  
hives facial swelling  
asthma (wheezing) shock  
skin reddening rash  
blisters

If an allergic reaction occurs, stop use and  
seek medical help right away.  
**Stomach bleeding warning:** This product  
contains an NSAID, which may cause severe  
stomach bleeding. The chance is higher if  
your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or  
steroid drug
- takes other drugs containing prescription  
or nonprescription NSAIDs (aspirin,  
ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs,  
except aspirin, increase the risk of heart  
attack, heart failure, and stroke. These can  
be fatal. The risk is higher if you use more  
than directed or for longer than directed.  
**Sore throat warning:** Severe or persistent  
sore throat or sore throat accompanied by high  
fever, headache, nausea, and vomiting may  
be serious. Consult doctor promptly. Do not  
use more than 2 days or administer to children  
under 3 years of age unless directed by doctor.

**Do not use**  
if the child has ever had an allergic  
reaction to ibuprofen or any other pain  
reliever/fever reducer



**Drug Facts (continued)**

- right before or after heart surgery
- Ask a doctor before use if**  
stomach bleeding warning applies to your child  
child has a history of stomach problems,  
such as heartburn  
child has problems or serious side effects  
from taking pain relievers or fever reducers  
child has not been drinking fluids  
child has lost a lot of fluid due to vomiting  
or diarrhea  
child has high blood pressure, heart disease,  
liver cirrhosis, kidney disease, or had a stroke  
child has asthma  
child is taking a diuretic

**Ask a doctor or pharmacist before use if  
the child is**  
under a doctor's care for any serious condition  
taking any other drug

**When using this product**  
give with food or milk if stomach upset occurs

- Stop use and ask a doctor if**
- child experiences any of the following signs  
of stomach bleeding:  
feels faint  
vomits blood  
has bloody or black stools  
has stomach pain that does not get better  
child has symptoms of heart problems or  
stroke:  
chest pain  
trouble breathing  
weakness in one part or side of body  
slurred speech  
leg swelling
  - the child does not get any relief within first  
day (24 hours) of treatment
  - fever or pain gets worse or lasts more than  
3 days
  - redness or swelling is present in the  
painful area
  - any new symptoms appear

**Keep out of reach of children.** In case of  
overdose, get medical help or contact a Poison  
Control Center right away. (1-800-222-1222)

**Directions**  
this product does not contain directions  
or complete warnings for adult use  
do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use  
weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not  
use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain  
child resistance
- wash dosage cup after each use

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\*or as directed by a doctor

**Other information**  
each 5 mL contains: sodium 2 mg  
do not use if printed neckband is  
broken or missing  
store at 20-25°C (68-77°F)  
do not freeze

**Inactive ingredients** anhydrous  
citric acid, D&C red #33, FD&C blue #1,  
FD&C red #40, glycerin, high fructose corn  
syrup, hypromellose, natural and artificial  
grape flavor, polysorbate 80, purified water,  
sodium benzoate, sorbitol solution,  
xanthan gum

**Questions or comments?**  
1-800-719-9260

**GLUTEN FREE**  
Distributed By  
AmerisourceBergen  
1300 Morris Drive  
Chesterbrook, PA 19087  
Questions or Concerns?  
[www.mygnp.com](http://www.mygnp.com)



13D526 29 C2

# GOOD NEIGHBOR PHARMACY CHILDRENS IBUPROFEN

ibuprofen suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46 122-110
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	PURPLE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-110-26	1 in 1 CARTON	04/02/2002	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA074937	04/02/2002	

