

**COLD AND FLU- acetaminophen, dextromethorphan hbr, phenylephrine
hcl solution**
Lidl US LLC

Lidl 44-012

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- liver disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

*Compare to the active ingredients in Vicks® DayQuil® Cold & Flu

NDC 71141-179-32

RELIEVES Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal Congestion, Cough

LIQUID

Daytime cold & flu RELIEF

Acetaminophen

Pain Reliever/Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Phenylephrine HCl

Nasal Decongestant

For ages 6 years
and over

**12 FL OZ
(355 mL)**

by **LIDL®**
F-012-02 ORG

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

*This product is not manufactured or distributed
by The Procter & Gamble Company, owner of the
registered trademark Vicks® DayQuil® Cold & Flu.
50844 ORG032301202

DISTRIBUTED BY
LIDL US, LLC
ARLINGTON, VA 22202
1-844-344-5071
www.lidl.com

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu
NDC 71141-179-32

RELIEVES Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal Congestion, Cough

Daytime
cold & flu
RELIEF
LIQUID

Acetaminophen
Pain Reliever/Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl

PEEL BACK FOR COMPLETE DRUG FACTS
TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms:
■ sore throat ■ nasal congestion
■ headache ■ minor aches and pains
■ cough due to minor throat and bronchial irritation
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if
■ adult takes more than 4,000 mg of acetaminophen in 24 hours
■ child takes more than 5 doses in 24 hours
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

DISTRIBUTED BY
LIDL US, LLC
ARLINGTON, VA 22202
1-844-344-5071
www.lidl.com

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Cold & Flu.
50844 ORG032301202

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Discard Seals, Empty & Replace Cap
5117817B
B-012-02 ORG
NPN20

For ages 6 years and over

For ages 6 years and over

Nasal Decongestant

12 FL OZ (355 mL)

by **LIDL**[®]
F-012-02 ORG

PLASTIC BOTTLE

4 056489 731849

no print / no varnish area
lot no. & exp. date

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease ■ diabetes ■ thyroid disease
- liver disease ■ high blood pressure

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ new symptoms occur

Drug Facts (continued)

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-800-426-9391

LIDL 44-012

COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71141-179
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71141-179-32	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/07/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/07/2023	

Labeler - Lidl US LLC (079389709)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(71141-179) , pack(71141-179)

Revised: 4/2025

Lidl US LLC