DICLOFENAC SODIUM- diclofenac sodium solution Advagen Pharma Ltd

Diclofenac Sodium Ophthalmic Solution, 0.1% Rx only Sterile Ophthalmic Solution

Prescribing Information

DESCRIPTION

Diclofenac sodium ophthalmic solution, 0.1% is a sterile, topical, nonsteroidal, antiinflammatory product for ophthalmic use. Diclofenac sodium is designated chemically as 2-[(2,6 dichlorophenyl) amino] benzeneacetic acid, monosodium salt, with an empirical formula of C ₁₄H ₁₀C _{l2}NO ₂Na. The structural formula of diclofenac sodium is:

Diclofenac sodium ophthalmic solution is available as a sterile solution, which contains diclofenac sodium 0.1%(1mg/mL).

*Inactive Ingredients:*polyoxyl 35 castor oil, Boric acid, tromethamine, sorbic acid (2 mg/mL), edetate disodium (1 mg/mL), and purified water.

Diclofenac sodium is a faintly yellow-white to light beige, slightly hygroscopic crystalline powder. It is freely soluble in methanol, sparingly soluble in water, very slightly soluble in acetonitrile, and insoluble in chloroform and in 0.1N hydrochloric acid. Its molecular weight is 318.14. Diclofenac sodium ophthalmic solution, 0.1% is a clear, colourless solution with an osmolality between 270 mOsmols/kg to 310 mOsmols/kg, buffered at a pH between 6.0 to 8.0. Diclofenac sodium ophthalmic solution has a faint characteristic odor of castor oil.

CLINICAL PHARMACOLOGY

Pharmacodynamics

Diclofenac sodium is one of a series of phenylacetic acids that has demonstrated antiinflammatory and analgesic properties in pharmacological studies. It is thought to inhibit the enzyme cyclooxygenase, which is essential in the biosynthesis of prostaglandins.

Animal Studies

Prostaglandins have been shown in many animal models to be mediators of certain kinds

of intraocular inflammation. In studies performed in animal eyes, prostaglandins have been shownto produce disruption of the blood-aqueous humor barrier, vasodilation, increased vascular permeability, leukocytosis, and increased intraocular pressure.

Pharmacokinetics

Results from a bioavailability study established that plasma levels of diclofenac following ocular instillation of two drops of Diclofenac Sodium Ophthalmic to each eye were below the limit of quantification (10 ng/mL) over a 4-hour period. This study suggests that limited, if any, systemic absorption occurs with Diclofenac Sodium Ophthalmic.

Clinical Trials

Postoperative Anti-Inflammatory Effects

In two double-masked, controlled, efficacy studies of postoperative inflammation, a total of 206 cataract patients were treated with Diclofenac Sodium Ophthalmic and 103 patients were treated with vehicle placebo. Diclofenac Sodium Ophthalmic was favored over vehicle placebo over a 2-week period for the clinical assessments of inflammation as measured by anterior chamber cells and flare.

In double-masked, controlled studies of corneal refractive surgery (radial keratotomy (RK) and laser photorefractive keratectomy (PRK)) patients were treated with Diclofenac Sodium Ophthalmic and/or vehicle placebo. The efficacy of Diclofenac Sodium Ophthalmic given before and shortly after surgery was favored over vehicle placebo during the 6-hour period following surgery for the clinical assessments of pain and photophobia. Patients were permitted to use a hydrogel soft contact lens with Diclofenac Sodium Ophthalmic for up to three days after PRK.

INDICATIONS AND USAGE

Diclofenac Sodium Ophthalmic is indicated for the treatment of postoperative inflammation in patients who have undergone cataract extraction and for the temporary relief of pain and photophobia in patients undergoing corneal refractive surgery.

CONTRAINDICATIONS

Diclofenac Sodium Ophthalmic is contraindicated in patients who are hypersensitive to any component of the medication.

WARNINGS

The refractive stability of patients undergoing corneal refractive procedures and treated with diclofenac sodium has not been established. Patients should be monitored for a year following use in this setting. With some nonsteroidal anti-inflammatory drugs, there exists the potential for increased bleeding time due to interference with thrombocyte aggregation. There have been reports that ocularly applied nonsteroidal antiinflammatory drugs may cause increased bleeding of ocular tissues (including hyphemas) in conjunction with ocular surgery.

There is the potential for cross-sensitivity to acetylsalicylic acid, phenylacetic acid derivatives, and other nonsteroidal anti-inflammatory agents. Therefore, caution should

be used when treating individuals who have previously exhibited sensitivities to these drugs.

PRECAUTIONS

General

All topical nonsteroidal anti-inflammatory drugs (NSAIDs) may slow or delay healing. Topical corticosteroids are also known to slow or delay healing. Concomitant use of topical NSAIDs and topical steroids may increase the potential for healing problems.

Use of topical NSAIDs may result in keratitis. In some susceptible patients continued use of topical NSAIDs may result in epithelial breakdown, corneal thinning, corneal infiltrates, corneal erosion, corneal ulceration, and corneal perforation. These events may be sight-threatening.

Patients with evidence of corneal epithelial breakdown should immediately discontinue use of topical NSAIDs and should be closely monitored for corneal health.

Postmarketing experience with topical NSAIDs suggests that patients experiencing complicated ocular surgeries, corneal denervation, corneal epithelial defects, diabetes mellitus, ocular surface disease (e.g., dry eye syndrome), rheumatoid arthritis, or repeat ocular surgeries within a short period-of time may be at increased risk for corneal adverse events, which may become sight- threatening. Topical NSAIDs should be used with caution in these patients.

Postmarketing experience with topical NSAIDs also suggests that use more than 24 hours prior to surgery or use beyond 14 days postsurgery may increase patient risk for occurrence and severity of corneal adverse events.

It is recommended that Diclofenac Sodium Ophthalmic, like other NSAIDs, be used with caution in patients with known bleeding tendencies or who are receiving other medications which may prolong bleeding time. Use of the same bottle for both eyes is not recommended with topical eye drops that are used in association with surgery.

Results from clinical studies indicate that Diclofenac Sodium Ophthalmic has no significant effect upon ocular pressure. However, elevations in intraocular pressure may occur following cataract surgery.

Information for Patients

Except for the use of a bandage hydrogel soft contact lens during the first 3 days following refractive surgery, Diclofenac Sodium Ophthalmic should not be used by patients currently wearing soft contact lenses due to adverse events that have occurred in other circumstances.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term carcinogenicity studies in rats given diclofenac sodium in oral doses up to 2 mg/kg/day (approximately 500 times the human topical ophthalmic dose) revealed no significant increases in tumor incidence. A 2-year carcinogenicity study conducted in mice employing oral diclofenac sodium up to 2 mg/kg/day did not reveal any oncogenic potential. Diclofenac sodium did not show mutagenic potential in various mutagenicity studies including the Ames test. Diclofenac sodium administered to male and female rats at 4 mg/kg/day (approximately 1000 times the human topical ophthalmic dose) did not

affect fertility.

Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger adult patients.

PREGNANCY

Teratogenic Effects

Reproduction studies performed in mice at oral doses up to 5,000 times (20 mg/kg/day) and in rats and rabbits at oral doses up to 2,500 times (10 mg/kg/day) the human topical dose have revealed no evidence of teratogenicity due to diclofenac sodium despite the induction of maternal toxicity and fetal toxicity. In rats, maternally toxic doses were associated with dystocia, prolonged gestation, reduced fetal weights and growth, and reduced fetal survival.

Diclofenac sodium has been shown to cross the placental barrier in mice and rats. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Non-teratogenic Effects

Because of the known effects of prostaglandin biosynthesis inhibiting drugs on the fetal cardiovascular system (closure of ductus arteriosus), the use of Diclofenac Sodium Ophthalmic during late pregnancy should be avoided.

Nursing Women

It is not known whether topical ophthalmic administration of Diclofenac Sodium Ophthalmic could result in sufficient systemic absorption to produce detectable quantities in breast milk. Because many drugs are excreted in human milk, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

ADVERSE REACTIONS

Ocular

Transient burning and stinging were reported in approximately 15% of patients across studies with the use of Diclofenac Sodium Ophthalmic. In cataract surgery studies, keratitis was reported in up to 28% of patients receiving Diclofenac Sodium Ophthalmic, although in many of these cases keratitis was initially noted prior to the initiation of treatment. Elevated intraocular pressure following cataract surgery was reported in approximately 15% of patients undergoing cataract surgery.

Lacrimation complaints were reported in approximately 30% of case studies undergoing incisional refractive surgery. The following adverse reactions were reported in approximately 10% or less of the patients: abnormal vision, acute elevated IOP, blurred

vision, conjunctivitis, corneal deposits, corneal edema, corneal opacity, corneal lesions, discharge, eyelid swelling, eye pain, injection (redness), iritis, irritation, itching, lacrimation disorder, and ocular allergy.

Systemic

The following adverse reactions were reported in 3% or less of the patients: abdominal pain, asthenia, chills, dizziness, facial edema, fever, headache, insomnia, nausea, pain, rhinitis, viral infection, and vomiting.

Clinical Practice

The following reactions have been identified during postmarketing use of topical diclofenac sodium ophthalmic solution, 0.1% in clinical practice. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. The reactions, which have been chosen for inclusion due to either their seriousness, frequency of reporting, possible causal connection to topical diclofenac sodium ophthalmic solution, 0.1%, or a combination of these factors, include corneal erosion, corneal infiltrates, corneal perforation, corneal thinning, corneal ulceration, and epithelial breakdown (see PRECAUTIONS, General).

To report SUSPECTED ADVERSE REACTIONS, contact Advagen Pharma Ltd, at 866-488-0312 or FDA at 1-800-FDA-1088 or *www.fda.gov/medwatch*.

OVERDOSAGE

Overdosage will not ordinarily cause acute problems. If Diclofenac Sodium Ophthalmic is accidentally ingested, fluids should be taken to dilute the medication.

DOSAGE AND ADMINISTRATION

Cataract Surgery

One drop of Diclofenac Sodium Ophthalmic should be applied to the affected eye, 4 times daily beginning 24 hours after cataract surgery and continuing throughout the first 2 weeks of the postoperative period.

Corneal Refractive Surgery

One or two drops of Diclofenac Sodium Ophthalmic should be applied to the operative eye within the hour prior to corneal refractive surgery. Within 15 minutes after surgery, one or two drops should be applied to the operative eye and continued 4 times daily for up to 3 days.

HOW SUPPLIED

Diclofenac Sodium Ophthalmic 0.1% (1 mg/mL) Sterile Solution is supplied in a white opaque LDPE bottle with a white opaque LDPE nozzle and grey coloured HDPE closure. The 2.5 mL fill is supplied in a 5 mL size bottle. The 5 mL fill is supplied in a 10 mL size bottle.

2.5 mL - **NDC**72888-150-21

5 mL - **NDC**72888-150-22

Store at 15°C to 25°C (59° to 77°F).

Dispense in original, unopened container only.

Distributed by:

Advagen Pharma Ltd.,

East Windsor, NJ 08520, USA.

Rev. 01/2025

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Diclofenac Sodium Ophthalmic 0.1% (1 mg/mL)- NDC 72888-150-21 - 2.5 mL Container label

5 0 2 1 5	FOR TOPICAL APPLICATION IN THE EYE Each mL contains 1 mg diclofenac sodium.	NDC 72888 Diclofena Ophthalmi	c Sodium c Solution	Distributed by: Advagen Pharma East Windsor, NJ, Code No.: MH/DR Rev. 03. 10/24	Ltd., 08520, USA UGS/28-KD/74 PLB0255	
N 728881	DOSAGE: See package insert. Store at 15°-25°C (59°-77°F).	Sterile Ophtha Rx Only	% Imic Solution 2.5 mL	101.00, 10,21	Overprinting Area 18.6 mm X 8 mm	

Diclofenac Sodium Ophthalmic 0.1% (1 mg/mL)- NDC 72888-150-21 - 2.5 mL Carton



Diclofenac Sodium Ophthalmic 0.1% (1 mg/mL)- NDC 72888-150-22 - 5 mL Container label



Diclofenac Sodium Ophthalmic 0.1% (1 mg/mL)- NDC 72888-150-22 - 5 mL Carton



DICLOFENAC SODIUM

diclofenac sodium solution

Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source)

NDC:72888-150

Route of Administration	OPHTHAL
-------------------------	---------

MIC **Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength DICLOFENAC SODIUM (UNII: QTG126297Q) (DICLOFENAC - UNII:14408QL0L1) DICLOFENAC SODIUM 1 mg in 1 mL **Inactive Ingredients Ingredient Name** Strength POLYOXYL 35 CASTOR OIL (UNII: 6D4M1DAL6O) BORIC ACID (UNII: R57ZHV85D4) **TROMETHAMINE** (UNII: 023C2WHX2V) SORBIC ACID (UNII: X045WJ989B) EDETATE DISODIUM (UNII: 7FLD91C86K) WATER (UNII: 059QF0K00R) Packaging Marketing Start **Marketing End** # Item Code **Package Description** Date Date **1** NDC:72888-150-21 1 in 1 CARTON 04/30/2025 2.5 mL in 1 BOTTLE; Type 0: Not a Combination 1

2	NDC:72888-150- 22	1 in 1 CARTON	04/30/2025					
2		5 mL in 1 BOTTLE; Type 0: Not a Combination Product						
Marketing Information								
M	larketing	Information						
M	larketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				

Labeler - Advagen Pharma Ltd (051627256)

Revised: 4/2025

Advagen Pharma Ltd