ACID REDUCER- ranitidine tablet McKesson

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do Not Use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with **lightheadedness**, **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° 25° C (68° 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

call 1-800-406-7984

PRINCIPAL DISPLAY PANEL

Health Mart_® **PHARMACY**

NDC 62011-0227-1

compare to Zantac 150® active ingredient[†]

ranitidine tablets, USP 150 mg

acid reducer

prevents and relieves heartburn associated with acid indigestion and sour stomach

maximum strength

24 tablets

Distributed by McKesson

5104604/0813





Other information

Gass or water symptoms, swallow 1 tablet with a glass or water of chinking betweeped by the order of chinking betweeped to the order of the order of chinking betweeped the cause hearburn ood of chinking betweeped the cause hearburn on one of chinking betweeped to the order of chinking betweeped to the order of chinking the order

Directions

a dults and children 12 years and over:

to relieve symptoms, swallow 1 tablet with a glass of water

a to prevent sumptoms, swallow 1 tablet with a

professional before use.

Reep out of reach of children. In case of overdog medical help or contact a Polson Control Cenight away (1-900-222-1222). If pregnant or breast-feeding, ask a health

Stop use and ask adotor it.

Stop use and ask adotor it.

Stop use and use this product for more than 14 days.

If days. Drug Facts (continued)

not use trouble or pain swallowing food, you have trouble or pain swallowing food, or black stoo condition.

Warnings

Allergy alert: Do not use if you are allergic to remitteline or other acid reducers

relieves heartburn associated with acid indigestion and sour stomach grevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and sour stomach brought on by certain foods were transes.

Agm 021 aniidiine P Ras raniidiine hydrochloride, USU 981 m Active ingredient (in each tablet) Purpose

Drug Facts



ACID REDUCER

ranitidine tablet

Contains

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:62011-0227

Route of Administration ORAL

Active Ingredient/Active Moiety

Basis of Strength Strength **Ingredient Name** RANITIDINE HYDRO CHLO RIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7) RANITIDINE 150 mg

Inactive Ingredients Ingredient Name Strength SILICON DIO XIDE (UNII: ETJ7Z6XBU4) CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48) HYPROMELLOSES (UNII: 3NXW29V3WO) FERRIC OXIDE RED (UNII: 1K09F3G675) MAGNESIUM STEARATE (UNII: 70097M6I30) CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U) POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) TALC (UNII: 7SEV7J4R1U) TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)

Product Characteristics Color Score no score OVAL (Oval Shaped) Shape Size 12mm Flavor 9 R Imprint Code

Packaging Item Code Package Description **Marketing Start Date Marketing End Date** 1 NDC:62011-0227-1 1 in 1 CARTON

1	24 in 1 BOTTLE					
Marketing Information						
Marketing Category	Application Number or Monograph Citation	n Marketing Start Date	Marketing End Date			
ANDA	ANDA200536	0 1/0 7/20 14				

Labeler - McKesson (177667227)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment					
Name	Address	ID/FEI	Business Operations		
Shasun Pharmaceuticals Limited		915786829	manufacture(62011-0227)		

Revised: 1/2014 McKesson