

**FIBER LAX- calcium polycarbophil tablet, film coated**  
**PD-Rx Pharmaceuticals, Inc.**

-----

**Drug Facts**

**Active ingredient (in each captab)**

Calcium polycarbophil 625 mg

**Purpose**

Bulk-Forming Laxative

(equivalent to polycarbophil 500 mg)

**Uses**

- relieves occasional constipation
- generally produces bowel movement in 12 to 72 hours

**Warnings**

**Choking:** Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

**Do not use**

longer than one week

**Ask a doctor before use if you have**

- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that lasts over two weeks

**Ask a doctor or pharmacist before use if you are**

taking any other drug. Take Fiber-Lax two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if**

- you have rectal bleeding
- you fail to have a bowel movement after using this product.

These may indicate a serious condition.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

Take Fiber-Lax with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.

- do not take more than 4 doses in 24 hours
- **adults & children 12 years and over:** 2 captabs 1 to 4 times a day
- **children under 12 years:** ask a doctor

## **Other information**

- **each captab contains:** calcium 140 mg
- store below 30°C (86°F) • do not refrigerate • protect from humidity

## **Inactive ingredients**

Caramel, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl sulfate and stearic acid

## ***Questions or comments?***

**Call 1-800-645-2158**

**TAMPER EVIDENT: DO NOT USE IF BLISTER SEAL IS BROKEN OR DAMAGED**

\*Rugby Laboratories, Inc. is not affiliated with the owner of the trademark FiberCon®.

Distributed by: **Rugby Laboratories**

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152


[www. Rugbylaboratories.com](http://www.Rugbylaboratories.com)

## **16 HOW SUPPLIED**

Fiber-Lax Tablets are available as follows:

500 mg: Off White, oval, scored tablet, debossed "CPC 339". Available in bottles of 30 captabs (NDC 55289-949-30)

Fiber-Lax

Drug Facts		Drug Facts (continued)
<b>Active Ingredient</b> (in each captab) Calcium polycarbophil 625mg (equivalent to polycarbophil 500 mg)	<b>Purpose</b> Bulk-Forming Laxative	<b>Directions</b> Take Fiber-Lax with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning. • do not take more than 4 doses in 24 hours • <b>adults &amp; children 12 years and over:</b> 2 captabs 1 to 4 times a day • <b>children under 12 years:</b> ask a doctor
<b>USES</b> • relieves occasional constipation • generally produces bowel movement in 12 to 72 hours		<b>Other information</b> • each captab contains: calcium 140mg • store below 30°C (86°F) • do not refrigerate • protect from humidity
<b>Warnings</b> <b>Choking:</b> Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.		<b>Inactive Ingredients</b> caramel, crospovidone, hypromellose, magnesium silicate, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silica, sodium lauryl sulfate and stearic acid
<b>Do not use</b> longer than one week		<b>Questions or comments?</b> 1-800-645-2158
<b>Ask a doctor before use if you have</b> • abdominal pain, nausea, or vomiting • noticed a sudden change in bowel habits that lasts over two weeks		 GTIN: 00355289949305 SNO: K20C890005 EXP: 11/2022 LOT: K20C89
<b>Ask a doctor or pharmacist before use if you are</b> taking any other drug. Take Fiber-Lax two or more hours before or after other drugs. Laxatives may affect how other drugs work.		
<b>Stop use and ask a doctor if</b> • you have rectal bleeding • you fail to have a bowel movement after using this product. These may indicate a serious condition.		
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.		

**NDC 55289-949-30**



# Fiber-Lax

## Bulk-Forming Laxative

- Promotes Regularity
- Non-Habit Forming

**500 mg**



Marketed and Packaged By:  
PD-Rx Pharmaceuticals, Inc  
Oklahoma City, OK 73127  
1-405-942-3040 v.8.19.0

**30 Captabs**  
TAMPER EVIDENT: DO NOT USE IF  
SAFETY SEAL IS BROKEN OR MISSING FROM BOTTLE.

## FIBER LAX

calcium polycarbophil tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55289-949(NDC:0536-4306)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CALCIUM POLYCARBOPHIL</b> (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARAMEL</b> (UNII: T9D99G2B1R)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	

<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	white (Off-white)	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	CPC;339
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55289-949-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/27/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	09/01/2010	

**Labeler** - PD-Rx Pharmaceuticals, Inc. (156893695)

**Registrant** - PD-Rx Pharmaceuticals, Inc. (156893695)

### Establishment

Name	Address	ID/FEI	Business Operations
PD-Rx Pharmaceuticals, Inc.		156893695	repack(55289-949)

Revised: 10/2024

PD-Rx Pharmaceuticals, Inc.