

HYOSCYAMINE SULFATE- hyoscyamine sulfate tablet
KMR Pharmaceuticals LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Hyoscyamine Sulfate
Sustained Release Tablets

Rx Only


DESCRIPTION

Each sustained-release tablet for oral administration contains:

Hyoscyamine 0.375 mg
Sulfate

Hyoscyamine sulfate is one of the principal anticholinergic/antispasmodic components of belladonna alkaloids.

Hyoscyamine sulfate is Benzeneacetic acid, α -(hydroxymethyl)-, 8- methyl-8-azabicyclo[3.2.1]oct-3-yl ester, [3(S)-endo]-, sulfate (2:1), dihydrate.



INACTIVE INGREDIENTS

Inactive ingredients include: calcium phosphate dibasic, ethylcellulose, green dye, lactose monohydrate, magnesium stearate (veg.), microcrystalline cellulose, and stearic acid.

CLINICAL PHARMACOLOGY

Hyoscyamine has actions similar to those of atropine, but is more potent in both its central and

peripheral effects. This product inhibits gastrointestinal propulsive motility and decreases gastric acid secretions. This product controls excessive pharyngeal, tracheal, and bronchial secretion. Once absorbed, this product disperses rapidly in the blood and is distributed throughout the entire body. The majority of hyoscyamine sulfate is excreted in the urine unchanged within the first 12 hours and only traces of hyoscyamine sulfate are found in the breast milk.

INDICATIONS AND USAGE

This product may be used in functional intestinal disorders to reduce symptoms such as those seen in mild dysenteries and diverticulitis. It can also be used to control gastric secretion, visceral spasm and hypermotility in cystitis, pylorospasm and associated abdominal cramps. Along with appropriate analgesics, this product is indicated in symptomatic relief of biliary and renal colic and as a drying agent in the relief of symptoms of acute rhinitis. This product is effective as adjunctive therapy in the treatment of peptic ulcer and irritable bowel syndrome, acute enterocolitis and other functional gastrointestinal disorders.

CONTRAINDICATIONS

Glaucoma, obstructive uropathy, obstructive diseases of the gastrointestinal tract, paralytic ileum, intestinal atony of elderly or debilitated patients, unstable cardiovascular status, severe ulcerative colitis, toxic megacolon, myasthenia gravis, and myocardial ischemia. This product is not recommended for use in children under twelve years of age.

WARNINGS

Heat prostration can occur with drug use in the event of high environmental temperature. Diarrhea may be an early symptom of incomplete intestinal obstruction, especially in patients with ileostomy or colostomy; in this instance, treatment would be inappropriate and possibly harmful. This product may cause drowsiness or blurred vision. Patients taking this product should be warned not to engage in activities requiring mental alertness such as operating a motor vehicle or other machinery or to perform hazardous tasks while taking this drug.

PRECAUTIONS

General

Use caution in patients with hiatal hernia associated with reflex esophagitis. Use extreme caution and only when needed in patients with autonomic neuropathy, hyperthyroidism, coronary heart disease, congestive heart failure and cardiac arrhythmia. Investigate any tachycardia before giving any anticholinergic drugs since they may increase the heart rate.

Prolonged use of anticholinergics may decrease or inhibit salivary flow, thus contributing to the development of caries, periodontal disease, oral candidiasis, and discomfort.

Information for Patients

The formulation for Hyoscyamine Sulfate Sustained Release Tablets is a non-eroding, non-gelling sustained release matrix. The drug releases from the tablet while the tablet remains intact. Therefore, it is possible that a tablet may appear in your stool.

This medication should be taken 30 minutes to one hour before meals. Tablets should be swallowed whole.

This medication should be used with caution during exercise or hot weather; overheating may result in heat stroke.

Hyoscyamine may cause drowsiness, dizziness or blurred vision; patients should observe caution before driving, using machinery or performing other tasks requiring mental alertness.

Drug Interactions

Absorption of other oral medications may be decreased during concurrent use with anticholinergics due to decreased gastrointestinal motility and delayed gastric emptying.

Drug interactions may occur when anticholinergics are used with the following medications: antacids, antidiarrheals (adsorbent), other anticholinergics, antimusculars, cyclopropane, haloperidol, ketoconazole, metoclopramide, opioid (narcotic) analgesics, and potassium chloride.

Pregnancy

Pregnancy Category C

Animal reproduction studies have not been conducted with this product. It is also not known whether this product can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Hyoscyamine crosses the placenta. This product should be given to a pregnant woman only if clearly needed.

Nursing Mothers

This product is excreted in human milk. This product should not be administered to a nursing mother.

Pediatric Use

This product is not recommended for use in children under twelve years of age. Infants and young children are especially susceptible to the toxic effects of anticholinergics. Close supervision is recommended for infants and children with spastic paralysis or brain damage since an increased response to anticholinergics has been reported in these patients and dosage adjustments are often required. When anticholinergics are given to children where the environmental temperature is high, there is a risk of a rapid increase in body temperature because of these medications' suppression of sweat gland activity.

A paradoxical reaction characterized by hyperexcitability may occur in children taking large doses of anticholinergics.

Geriatric Use

Geriatric patients may respond to usual doses of anticholinergics with excitement, agitation, drowsiness, or confusion.

Geriatric patients are especially susceptible to the anticholinergic side effects, such as constipation, dryness of mouth, and urinary retention (especially in males). If these side effects occur and continue or are severe, medication should probably be discontinued.

Caution is also recommended when anticholinergics are given to geriatric patients, because of the danger of precipitating undiagnosed glaucoma.

Memory may become severely impaired in geriatric patients, especially those who already have memory problems, with the continued use of anticholinergics since these drugs block the actions of acetylcholine, which is responsible for many functions of the brain, including memory functions.

ADVERSE REACTIONS

Not all of the following adverse reactions have been reported with hyoscyamine sulfate. The following adverse reactions have been reported for pharmacologically similar drugs with anticholinergic-antispasmodic action. Adverse reactions may include dryness of the mouth, urinary hesitancy and

retention; blurred vision; tachycardia; palpitations; mydriasis; cycloplegia; increased ocular tension; loss of taste; headache; nervousness; drowsiness; weakness; dizziness; insomnia; nausea; vomiting; impotence; suppression of lactation; constipation; bloated feeling; allergic reactions or drug idiosyncrasies; urticaria and other dermal manifestations; ataxia; speech disturbance; some degree of mental confusion and/or excitement (especially in elderly persons); and decreased sweating.

OVERDOSAGE

The signs and symptoms of overdose are headache, nausea, vomiting, blurred vision, dilated pupils, hot dry skin, dizziness, dryness of the mouth, difficulty in swallowing and CNS stimulation. Measures to be taken are immediate lavage of the stomach and injection of physostigmine 0.5 to 2 mg intravenously and repeated as necessary up to a total of 5 mg. Fever may be treated symptomatically (tepid water sponge baths, hypothermic blanket). Excitement to a degree which demands attention may be managed with sodium thiopental 2% solution given slowly intravenously or chloral hydrate (100-200 mL of a 2% solution) by rectal infusion. In the event of progression of the curare-like effect to paralysis of the respiratory muscles, artificial respiration should be instituted and maintained until effective respiratory action returns.

In rats, the LD₅₀ for hyoscyamine is 375 mg/kg. Hyoscyamine is dialyzable.

DOSAGE AND ADMINISTRATION

Adults and children 12 years or older: 1 or 2 tablets every 12 hours. Dosage may be adjusted to 1 tablet every 8 hours if needed. Do not exceed 4 tablets in 24 hours. Tablets should be swallowed whole.

Note: Geriatric patients may be more sensitive to the effects of the usual adult dose.

HOW SUPPLIED

Hyoscyamine Sulfate Sustained Release Tablets are supplied as green, capsule shaped tablets debossed "203" on one side and scored on the opposite side. Available in bottles of 100 tablets, NDC 57629-203-10.

Storage and Handling

Dispense in a tight, light-resistant container as defined in USP/NF, with a child-resistant closure. Store at controlled room temperature between 20°-25°C (68°- 77°F), see USP Controlled Room Temperature.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN. IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY.

Distributed by:

KMR Pharmaceuticals LLC
New York, NY 10022

PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label

NDC 57629-203-10


**Hyoscyamine
Sulfate**

Sustained Release Tablets

Each sustained-release tablet contains:
Hyoscyamine Sulfate 0.375 mg

Rx Only
100 Tablets

KMR Pharmaceuticals LLC

Lot No. / Exp. Date		USUAL DOSAGE: <i>Adults and pediatric patients 12 years of age and older: 1 to 2 tablets every 12 hours. See attached insert for further information.</i>	NDC 57629-203-10
		Store at controlled room temperature between 20°- 25°C (68°- 77°F), see USP Controlled Room Temperature. Dispense in tight, light-resistant containers as defined in USP/NF with a child-resistant closure. Distributed by: KMR Pharmaceuticals LLC New York, NY 10022 400824-03 Iss. 01/2014	<h1 style="color: red;">Hyoscyamine Sulfate</h1> <h2 style="color: red;">Sustained Release Tablets</h2> <p>Each sustained-release tablet contains: Hyoscyamine Sulfate 0.375 mg</p> <p style="text-align: center;">Rx Only 100 Tablets</p> <h2 style="color: red;">KMR Pharmaceuticals LLC</h2>

HYOSCYAMINE SULFATE

hyoscyamine sulfate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:57629-203
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hyoscyamine Sulfate (UNII: F2R8V82B84) (Hyoscyamine - UNII:PX44XO846X)	Hyoscyamine Sulfate	0.375 mg

Inactive Ingredients

Ingredient Name	Strength
calcium phosphate, dibasic, anhydrous (UNII: L11K75P92J)	
ethylcelluloses (UNII: 7Z8S9VYZ4B)	
lactose monohydrate (UNII: EWQ57Q8IX)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
stearic acid (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	14mm
Flavor		Imprint Code	203
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57629-203-10	100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		01/01/2011	

Labeler - KMR Pharmaceuticals LLC (078628750)

Revised: 7/2015

KMR Pharmaceuticals LLC