CAREONE TUSSIN CF MULTI SYMPTOM COLD- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Tussin CF Multi-Symptom Cold Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredients in Robitussin® Multi-Symptom Cold

TUSSIN CF MULTI-SYMPTOM COLD

Cough Suppressant-Dextromethorphan HBr

Expectorant-Guaifenesin

Nasal Decongestant-Phenylephrine HCl
Relieves:
Cough
Nasal Congestion
Mucus
Adult
For Ages 12 & Over
Non-Drowsy
Alcohol Free
Gluten Free
OUR PHARMACISTS RECOMMEND
Cherry Flavor
4 FL OZ (118mL)



Mucus

Non-Drowsy

Alcohol Free

Gluten Free

For Ages 12 & Over

ompare to the active ingredients in Robitussin® Multi-Symptom Cold*

Cough Suppressant-Dextromethorphan HBr Expectorant-Gualfenesin Nasal Decongestant-Phenylephrine HCI

- · Cough
- Nasal Congestion

CARE**ONE**®

• Mucus

For Ages 12 & Over

Non-Drowsy

Alcohol Free



4 FL OZ (118mL)

Drug Facts

Active ingredients Purposes (in each 10 mL)
Dextromethorphan HBr,

..Cough suppressant

IS BROKEN OR MISSING
DO NOT USE IF PRINTED NECKBAND

...Nasal decongestan

- Uses helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes temporarily relieves these symptoms
- occurring with a cold:

 nasal congestion
 cough due to minor throat and bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
 high blood pressure
 thyroid disease
 diabetes
- tryroid disease diabetes

 I trouble urinating due to an enlarged
 prostate gland

 cough that occurs with too much
 phlegm (mucus)

 cough that lasts or is chronic such as occurs
 with assigning authors observed hereals in
- with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
 symptoms do not get better within 7 days
 or are accompanied by fever
 cough lasts for more than 7 days, comes
- back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or cor Poison Control Center right away

- Directions

 do not take more than 6 doses in any 24-hour period

 measure only with dosing cup provided

 keep dosing cup with product

 null = milliliter

- this adult product is not intended for use

in children under 12 years of age		
age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

- sodium 5 ma
- Other information
 each 10 mL contains: sod
 store at 20-25°C (68-77°F).
 Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Multi-Symptom Cold. DISTRIBUTED BY FOODHOLD U.S.A., LLC LANDOVER, MD 20785 · 1-877-846-9949 ©2017 S&S Brands, LLC Quality guaranteed or your money back.

PARENTS: Learn about teen medicine abuse

www.StopMedicineAbuse.org



Active Ingredient/Active Moiety

Cherry Flavor



LOT NO.

EXP.

: 51626 OF C1

CAREONE TUSSIN CF MULTI SYMPTOM COLD

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information HUMAN OTC DRUG NDC:41520-937 Product Type Item Code (Source) ORAL Route of Administration

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
EDETATE DISO DIUM (UNII: 7FLD9 1C86K)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		

Product Characteristics			
Color	RED (clear)	Score	
Shape		Size	
Flavor	FRUIT (Menthol)	Imprint Code	
Contains			

I	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:41520-937-26	1 in 1 CARTON	08/05/2016		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/05/2016	

Labeler - American Sales Company (809183973)

Revised: 8/2017 American Sales Company