

**ASPIRIN LOW DOSE- aspirin tablet, delayed release  
DOLGENCORP, LLC**

-----  
**Dollar General 44-645**

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

**Questions or comments?**

**1-888-309-9030**

**Principal display panel**

**DG™** | health

**Compare to active  
ingredient of St. Joseph®  
Low Dose Aspirin\*\***

**Low Dose  
Aspirin**

Pain Reliever (**NSAID**)

- Enteric coated
- Aspirin regimen

**81 mg  
each**

Talk to your doctor or  
other healthcare provider  
before using this product  
for your heart.

**100 Tablets**

Actual Tablet Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER IS BROKEN OR MISSING**

This product is not manufactured or distributed by  
Foundation Consumer Healthcare, LLC, owner of the  
registered trademark St. Joseph® Low Dose Aspirin.

50844

REV1024D64512

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**100%  
Satisfaction  
Guaranteed!**  
(888) 309-9030

No Print / No Varnish  
Lot no. & Exp. date

A0532

N  
3  
50844 41412  
0

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Compare to active ingredient of St. Joseph® Low Dose Aspirin\*\*

Low Dose

# Aspirin

11 mg each

Pain Reliever (NSAID)

- Enteric coated
- Aspirin regimen

100 Tablets

Actual Tablet Size

100% Satisfaction Guaranteed!

(888) 309-9030

\*\*This product is not manufactured or distributed by Foundation Consumer Healthcare, LLC, owner of the registered trademark St. Joseph® Low Dose Aspirin. 50844 REV1024D64512

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

B-0315-645-12-RR  
REV1024D64512

**Drug Facts (continued)**

unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, trehalyl citrate

**Questions or comments?**  
1-888-309-9030

**Drug Facts (continued)**

- have 3 or more alcoholic drinks every day while using this product.

**Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy**

A0532

N  
3  
50844 41412  
0

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts (continued)**

unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, trehalyl citrate

**Questions or comments?**  
1-888-309-9030

**Drug Facts (continued)**

- have 3 or more alcoholic drinks every day while using this product.

**Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts (continued)**

unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, trehalyl citrate

**Questions or comments?**  
1-888-309-9030

**Drug Facts (continued)**

- have 3 or more alcoholic drinks every day while using this product.

**Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy**

TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Dollar General 44-645

<p style="font-weight: bold; font-size: 1.2em; margin: 0;">ASPIRIN LOW DOSE</p> <p style="margin: 0;">aspirin tablet, delayed release</p>			
<p style="font-weight: bold; margin: 0;">Product Information</p>			
<p style="font-weight: bold; margin: 0;">Product Type</p>	<p>HUMAN OTC DRUG</p>	<p style="font-weight: bold; margin: 0;">Item Code (Source)</p>	<p>NDC:55910-945</p>
<p style="font-weight: bold; margin: 0;">Route of Administration</p>	<p>ORAL</p>		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	L
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-945-12	1 in 1 CARTON	07/25/2014	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/25/2014	

**Labeler** - DOLGENCORP, LLC (068331990)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-945)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-945)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-945)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55910-945)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55910-945)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-945)

Revised: 8/2025

DOLGENCORP, LLC