

**HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin and caffeine tablet, film coated**  
**Grocery Outlet**

-----  
**Headache Relief**

**Active ingredient(s)**

Acetaminophen 250 mg  
Aspirin 250 mg (NSAID\*)  
Caffeine 65 mg

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever  
Pain reliever  
Pain reliever aid

**Use(s)**

temporarily relieves minor aches and pains due to

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling

- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 8 tablets in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

#### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

#### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

#### **Ask a doctor or pharmacist before use if**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or under a doctor's care for any serious condition

#### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood

- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- ever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition

### **If pregnant or breastfeeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice the signs or symptoms.

### **Directions**

- do not use more than directed
- drink a full glass of water with each dose.

**adults and children 12 years and over:** take 2 tablets every 6 hours; not more than 8 tablets in 24 hours

**children under 12 years:** ask a doctor

### **Other information**

store at 20°-25°C (68°-77°F)

### **Inactive ingredients**

corn starch, croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

### **Questions/Comments**

Call **1-877-290-4008**

### **Principal Display Panel**

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

LE2416 R1175

PRODUCT OF USA

www.grocerypartner.com

Emeryville, CA 94608

Grocery Outlet, Inc.

DISTRIBUTED BY:

Extra Strength

US Holdings LLC, owner of the registered trademark Excedrin®

100% MONEY BACK GUARANTEE

COMBINATION CAP

**Drug Facts (continued)**

The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat

**Do not use** ■ if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are** ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of

**Drug Facts (continued)**

hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice the signs or symptoms.

**Directions** ■ do not use more than directed ■ drink a full glass of water with each dose.

■ adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours

■ children under 12 years: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

**Inactive ingredients** corn starch, croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

**Questions or comments?** Call 1-877-290-4008

NDC 85829-285-93

Compare to active ingredients in Excedrin® Extra Strength\*\*

**EXTRA STRENGTH**

**Headache Relief**

**ACETAMINOPHEN, 250 mg ASPIRIN, 250 mg (NSAID\*)**

**CAFFEINE, 65 mg**

Pain Reliever / Pain Reliever Aid



**Drug Facts**

Active ingredients (in each tablet)	Purposes
Acetaminophen 250mg	Pain reliever
Aspirin 250mg (NSAID*)	Pain reliever
Caffeine 65mg	Pain reliever aid
*nonsteroidal anti-inflammatory drug	

**Uses** temporarily relieves minor aches and pains due to ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache ■ premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert: Aspirin** may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 tablets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding.



Lot: UNWARNISH  
Exp:

Carton

NDC 85828-285-93

**EXTRA STRENGTH**  
**Headache Relief**  
**ACETAMINOPHEN, 250 mg**  
**ASPIRIN, 250 mg (NSAID)**  
**CAFFEINE, 65 mg**  
 Pain Reliever / Pain Reliever Aid

**GO** **24 TABLETS**

**SEE CARTON FOR COMPLETE PRODUCT INFORMATION**  
**DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.  
**Allergy alert:** Acetaminophen may cause severe skin reactions.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
**Liver warning:** This product contains acetaminophen.  
**Stomach bleeding warning:** This product contains an NSAID which may cause severe stomach bleeding.  
**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat  
**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**  
**Directions** ■ do not use more than directed ■ drink a full glass of water with each dose.  
 ■ adults and children 12 years and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours  
 ■ children under 12 years: ask a doctor

**100% MONEY BACK GUARANTEE**

**DISTRIBUTED BY:**  
 Grocery Outlet, Inc.  
 Emeryville, CA 94608  
 www.groceryoutlet.com

PRODUCT OF USA. LB2415 R1125

Lot.:  
 Exp.:  
**UNVARNISH**

Label

## HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin and caffeine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:85828-285
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY35J)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
--------------	-------	--------------	----------

<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ 285
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-285-93	1 in 1 CARTON	03/17/2026	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/17/2026	

**Labeler** - Grocery Outlet (029161585)