STOMACH RELIEF- bismuth subsalicylate capsule, liquid filled Spirit Pharmaceuticals LLC

Stomach Relief

Drug Facts

Active Ingredient (in each tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and anti-diarrheal

Uses

relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate.

Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products

Do not use if you have

• an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have

• fever • mucus in the stool

Ask a doctor or pharmacist before use if you are if you are

you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product

• symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

• swallow with water, do not chew • adults and children 12 years and over: • 2 softgels (1 dose) every 1/2 hour or 4 softgels (2 doses) every hour as needed for diarrhea • 2 soft gels (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) • do not exceed 8 doses (16 softgels) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

• each softgel contains: Salicylate 99 mg • very low sodium • store between 15–30°C (59–86°F)

Inactive ingredients

Aerosil, beeswax, FD&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide

Questions or comments?

1-888-333-9792

Principal Display Panel





CABINET: Stomach Relief Refill

COMPARE TO THE ACTIVE INGREDIENT IN-

Pepto-Bismol®*

Bismuth Subsalicylate 262 mg

DEL LEVEE

NAUSEA · HEARTBURN · INDIGESTION · UPSET STOMACH DIARRHEA



TO OPEN:

1. POSITION THUMBS INSIDE THE SHORT FLANGE OVER THE LOCKS. 2. GRIP WHILE PIVOTING THUMBS OUTWARD TO OPEN THE L. CKS.

Temper evident: do not use if pouch is open

Drug Facts

Active Ingredient (in each tablet)

Purpose

Bismuth subsalicylate

Upset stomach reliever 262 mg,..... and anti-diarrheal

Uses relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including:
• heartbum
• indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate.

Do not take if you are

allergic to salicylates (including aspirin)
 taking other salicylate

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have

fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

• symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions • swallow with water, do not chew • adults and children 12 years and over: • 2 softgels (1 dose) every 1/2 hour or 4 softgels (2 doses) every hour as needed for diarrhea • 2 softgels (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) • do not exceed 8 doses (16 softgels) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Drug Facts (continued)

Other Information • each softgel contains: Salicylate 99 mg • very low sodium • store between 15–30°C (59–86°F)

Inactive ingredients Aerosil, beeswax, FD&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide

Questions or comments? 1-888-333-9792

It takes guts to eliminate plastic.

- * This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Pepto-Bismol®
- † Less than the limit of detection and consistent with gluten-free diet labeling per FDA

ACC-1-10020 Distributed by: Cabinet Health P.B.C. Question/Comments? 1-888-333-9792 www.cabinethealth.com

Lot / Expiration -









STOMACH RELIEF

bismuth subsalicylate capsule, liquid filled

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:68210-4238

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID -**BISMUTH** 262 mg **SUBSALICYLATE**

UNII:O414PZ4LPZ)

Inactive Ingredients Ingredient Name Strength SYNTHETIC BEESWAX (UNII: 08MNR5YE2R) SILICON (UNII: Z4152N8IUI) D&C RED NO. 33 (UNII: 9DBA0SBB0L) FD&C YELLOW NO. 6 (UNII: H77VEI93A8) GELATIN (UNII: 2G86QN327L) **GLYCERIN** (UNII: PDC6A3C0OX) PARAFFIN (UNII: 1900E3H2ZE) WATER (UNII: 059QF0KO0R) LECITHIN, SOYBEAN (UNII: 1DI56QDM62) HYDROXYMETHYL CELLULOSE (UNII: 273FM27VK1) **SORBITOL SOLUTION** (UNII: 8KW3E207O2)

Product Characteristics				
Color	pink	Score	no score	
Shape	CAPSULE	Size	14mm	
Flavor		Imprint Code	262	
Contains				

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
NDC:68210- 4238-5	60 in 1 POUCH; Type 0: Not a Combination Product	11/17/2022		

Marketing	Information

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Marketing Category

Application Number or Monograph Citation

Marketing Start Date

Marketing End Date

Labeler - Spirit Pharmaceuticals LLC (179621011)

Revised: 12/2024 Spirit Pharmaceuticals LLC