

NIGHTTIME COLD AND FLU MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled
Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Safeway Nighttime Cold & Flu

Active Ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan Hydrobromide 15 mg

Doxylamine Succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

Other Information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive Ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, titanium dioxide*, white ink

*may contain this ingredient

Questions or Comments?

Call **1-877-753-3929 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®†

Nighttime Liquid Caps

Multi-Symptom Cold & Flu Relief

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

Pain reliever / Fever reducer

Cough suppressant

Antihistamine

Alcohol-free

Softgels** (**Liquid-Filled Capsules)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble

Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Another Quality Product

Distributed by McKesson

One Post Street, San Francisco, CA 94104

www.healthmart.com/healthmartbrand

Package Label



Health Mart

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®†

NDC 62011-0389-1

Nighttime Liquid Caps

Multi-Symptom Cold & Flu Relief

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Pain reliever/Fever reducer
Cough suppressant
Antihistamine



ACTUAL SIZE

PER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

No.:
Date:

<p>Drug Facts (continued)</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves common cold and flu symptoms sore throat headache minor aches and pains runny nose and sneezing fever cough due to minor throat and bronchial irritation 	<p>Drug Facts</p> <p>Active ingredients (in each softgel)</p> <ul style="list-style-type: none"> Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 15 mg.....Cough suppressant Doxylamine succinate 6.25 mg.....Antihistamine
---	--

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed (see Overdose warning) adults and children 12 years and over: take 2 softgels with water every 6 hours swallow whole; do not crush, chew, or dissolve children under 12 years: do not use when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing <p>Other information</p> <ul style="list-style-type: none"> store between 15-30°C (59-86°F) avoid excessive heat <p>Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide, white ink</p> <p>*may contain this ingredient</p> <p>Questions or comments?</p> <p>Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours</p> <ul style="list-style-type: none"> with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions.</p> <ul style="list-style-type: none"> skin reddening blisters rash <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease glaucoma trouble urinating due to an enlarged prostate gland a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers <p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage excitability may occur, especially in children avoid alcoholic drinks marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery
--	---

16 Softgels** (**Liquid-Filled Capsules)

Alcohol-free



Lot Exp

McKesson

Another Quality Product
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
healthmart.com

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



PLD-E41T FC005090

HEALTH MART Multi-Symptom Cold & Flu Relief

NIGHTTIME COLD AND FLU MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0389
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	P30;94A;AP017
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0389-1	16 in 1 CARTON	09/30/2018	05/31/2024
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2018	05/31/2024

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 9/2022

Strategic Sourcing Services LLC