

**HEALTH SENSE INFANTS IBUPROFEN ORAL SUSPENSION- ibuprofen suspension  
Dolgenercorp Inc**

-----  
**Dollar General Health Sense Infants' Ibuprofen Oral Suspension, Berry Flavor Drug Facts**

**Active ingredient (in each 1.25 mL)**

Ibuprofen 50 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding.

The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if the child has**

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- taken a diuretic

**Ask a doctor or pharmacist before use if the child is**

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor’s care for any serious condition
- taking any other drug

**When using this product**

- give with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

**Stop use and ask a doctor if**

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- shake well before using
- find the right dose on chart below. If possible, use weight to dose; otherwise use age.
- measure with dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child’s mouth, toward the inner cheek
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

## Other information

- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

## Inactive ingredients

artificial flavor, butylparaben, citric acid, FD&C red #40, glycerin, hypromellose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

## Questions or comments?

1-888-309-9030

## Principal Display Panel

Compare to active ingredient of Motrin® Infants' Drops

Concentrated

For Ages 6 mos.-23 mos.

Infants' Ibuprofen Oral Suspension

50 mg per 1.25 mL

Pain Reliever/Fever Reducer (NSAID)

Lasts up to 8 hours

Alcohol Free

Use Only With Enclosed Syringe

Berry Flavor



Product Carton



**Product Label**

**HEALTH SENSE INFANTS IBUPROFEN ORAL SUSPENSION**

ibuprofen suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-057
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10 QM) (IBUPROFEN - UNII:WK2XYI10 QM)	IBUPROFEN	50 mg in 1.25 mL

**Product Characteristics**

<b>Color</b>	PINK (light)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-057-05	1 in 1 CARTON		
1		15 mL in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075217	06/08/2009	

**Labeler** - Dolgencorp Inc (068331990)