

**IBUPROFEN- ibuprofen tablet
WALGREENS**

Walgreens Ibuprofen 200 mg Tablets - Brown Caplets

DRUG FACTS

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially

important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and over:	<ul style="list-style-type: none">• take 1 caplet every 4 to 6 hours while symptoms persist• if pain or fever does not respond to 1 caplet, 2 caplets may be used• do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years:	<ul style="list-style-type: none">• ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call **1-877-290-4008**

Varnish Omit

Lot No.:
Exp Date:



ITEM 720587 W000000-00000-0
MADE IN INDIA
walgreens.com ©2021 Walgreen Co.
100% SUSTAINABLE GUARANTEED
200 WILMOT RD., DEERFIELD, IL 60015
DISTRIBUTED BY: WALGREEN CO.
©2021 Walgreen Co. All rights reserved.
Walgreen is a registered trademark of Walgreen Co.

**READ ALL PRODUCT INFORMATION BEFORE USING.
KEEP THE BOX FOR IMPORTANT INFORMATION.**

Drug Facts

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID) *...Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
● headache ● toothache ● backache
● menstrual cramps ● the common cold
● muscle aches ● minor pain of arthritis
● temporarily reduces fever

Warnings

- Alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
● hives ● facial swelling ● asthma (wheezing) ● shock ● skin redness ● rash ● blisters
If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
● are age 60 or older
● have had stomach ulcers or bleeding problems
● take a blood-thinning (anticoagulant) or steroid drug
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
● have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
● if you have ever had an allergic reaction to any other pain reliever/fever reducer
● right before or after heart surgery

Ask a doctor before use if
● you have problems or serious side effects from taking pain relievers or fever reducers
● stomach bleeding warning applies to you
● you have a history of stomach problems, such as heartburn

● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
● you are taking a diuretic
Ask a doctor or pharmacist before use if you are
● under a doctor's care for any serious condition
● taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
● taking any other drug

When using this product:
● take with food or milk if stomach upset occurs
Stop use and ask a doctor if
● you experience any of the following signs of stomach bleeding:
● faint ● vomit ● blood ● have bloody or black stools
● have stomach pain that does not get better
● you have symptoms of heart problems or stroke:
● chest pain ● trouble breathing

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call 1-877-290-1008
TAP HERE TO GET THE FULL PRODUCT INFORMATION. THE BOTTOM IS CUT OPEN FOR EASY ACCESS.

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.



Drug Facts (continued)

- weakness in one part of side of body
- slurred speech ● big swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years:
● ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

W50RG0122-F
600R 1021
REV-0423

NDC 0363-9800-01



Compare to the active ingredient in Advil® Caplets*

Ibuprofen

TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

100 CAPLETS** (**capsule-shaped tablets)



ACTUAL SIZE

Varnish Omit

Varnish Omit

Varnish/Print/Omit Area

E827

Adhesive Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery.

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke ■ you are taking a diuretic.

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:

- pain, heartburn, indigestion, or heartburn
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

 ■ you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- big swelling

 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed

For the smallest effective dose should be used

adults and children 12 years and over:	<ul style="list-style-type: none"> ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years:	<ul style="list-style-type: none"> ■ ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68°-77°F)

Inactive ingredients calcium silicate, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call 1-877-296-4008

VALUE SIZE NDC 0383-9600-31

Walgreens

Compare to the active ingredient in Advil® Caplets*

Ibuprofen

TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

300 CAPLETS**

(**capsule-shaped tablets)



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR OTHERWISE DAMAGED

Drug Facts

Active ingredient (in each caplet) Ibuprofen 200 mg (NSAID)*

Purpose Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- skin redness/itching
- rash
- dizziness
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to ask a pharmacist for more information. This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Caplets.

ITEM 666907 W000000-0000-0
 DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com 1-800-955-1733
 ©2021 Walgreen Co.



3 111917 20453 6

W5ORG0122-F
800R 1101 REV1222 Made in India

Lot No.:

Varnish Omit Area

Exp Date:



Varnish Omit
Lot No.:
Exp. Date:



This product is not manufactured or distributed by Pfizer Consumer Healthcare's owner of the registered trademark KNOX® Caplets.



READ ALL INFORMATION BEFORE USING. KEEP THE BOX FOR PROPER STORAGE INFORMATION.

Drug Facts

Active ingredient (in each caplet): Purpose
Ibuprofen
200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains due to:
● headache ● toothache ● backache ● menstrual cramps ● the common cold ● muscle aches ● minor pain of arthritis ■ Temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ dizziness. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 65 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs combining prescription or over-the-counter NSAIDs (aspirin, naproxen, ibuprofen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver problems, kidney disease, asthma, or had a stroke ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product:
■ Take with food or milk in stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding: ● faintness ● vomit blood ● have bloody or black stools
■ have stomach pain that does not get better

Drug Facts (continued)

■ you have symptoms of heart problems or stroke:
● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech
● big swelling ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swelling is present in the genital area ● any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used
adults and children 12 years
■ take 1 caplet every 4 to 6 hours with symptoms present
■ If pain or fever does not respond to 1 caplet, 2 caplets may be used. ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years:
■ ask a doctor
Other information ■ read warnings and

Drug Facts (continued)
■ store at 20-25°C (68-77°F)
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide
Questions or comments? call 1-877-290-4008
IN THE DRUG STORE OR ONLINE AT WALGREENS.COM, THIS MAY OVER BE MONITORED AND RE-OPENED FOR THE FOLLOWING REASONS:
Your pharmacist will advise you of these brands.

Varnish Omit

Varnish Omit



Ibuprofen
TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

24 CAPLETS**
(**capsule-shaped tablets)



ACTUAL SIZE

926 E Varnish/Print Omit Area

W3ORG0122-F
800R 1021
REL40423

NDC 0363-9600-42

Adhesive Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist

■ if pain or fever does not respond to 1 caplet, 2 caplets may be used

■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Questions or comments? call 1-877-290-4008

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2021 Walgreen Co.
MADE IN INDIA 600R 1021
ITEM Z20789, W00000-0000-0.

W3ORG0122-F

NDC 0363-9600-42

Walgreens

Ibuprofen

TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

24 CAPLETS**
(**capsule-shaped tablets)

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THE BOX FOR IMPORTANT INFORMATION.

Uses ■ temporarily relieves minor aches and pains due to: ● headache ● toothache ● backache ● the menstrual cramps ● minor common cold ● muscular aches ● minor pain of arthritis ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued under label)

Lot No.: **Varnish Omit Area**

PEEL HERE

Exp Date:

NDC 0363-9600-01

Walgreens

Ibuprofen

TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

100 CAPLETS**
(**capsule-shaped tablets)

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: ● headache ● toothache ● backache ● the menstrual cramps ● minor common cold ● muscular aches ● minor pain of arthritis ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? call 1-877-290-4008

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2021 Walgreen Co.
MADE IN INDIA
W3ORG0122-F ITEM 720587
600R 1021 40114802 Code No.: W00000-0000-0
G0DRUGS/5/5

Lot No.: **Varnish Omit Area**

Exp Date:



Adhesive Area

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**
Directions ■ do not take more than directed ■ the smallest effective dose should be used
 adults and children 12 years and over:
 ■ take 1 caplet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
 ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
 children under 12 years: ■ ask a doctor
Questions or comments? call 1-877-290-4008
 DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 Walgreens.com ©2021 Walgreen Co.
 MADE IN INDIA 600 R 1021
 ITEM 718886 W00000-0000-0
 W3ORG0122-F

NDC 0363-9600-05

Walgreens

Ibuprofen

TABLETS USP, 200 mg /
 PAIN RELIEVER /
 FEVER REDUCER (NSAID)

50 CAPLETS**
 (**capsule-shaped tablets)

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: ● headache ● toothache ● backache ● menstrual cramps ● the common cold ● muscular aches ● minor pain of arthritis ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued under label)

40114702 Code No.: G0DRUGS/515

Varnish Omit Area

Lot No.:

Exp. Date:



IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9600
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	

FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	brown	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	117
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9600-42	24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
2	NDC:0363-9600-05	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
3	NDC:0363-9600-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
4	NDC:0363-9600-31	300 in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	12/11/2018	
5	NDC:0363-9600-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	12/11/2018	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-9600)